

Lesson 10 – Inverses & Inverse Functions

IBHL1 Math - Santowski

9/21/16

HL1 Math - Santowski

1

Fast Five – Skills Preview

- Isolate “x” in the following equations (make “x” the subject of the equation)

(a) $3x - 2y + 7 = 0$

(b) $f(x) = \frac{1}{2}(x+2)^2 - 5$

(c) $h(x) = \frac{2}{3-x}$

(d) $y = x^2 - 4x + 1$

(e) $y = e^{x+1}$

9/21/16


HL1 Math - Santowski

2

Fast Five – Concept Preview

Determine:

- (a) Domain of $f(x)$
- (b) Range of $f(x)$
- (c) Mapping of inverse
- (d) Domain of inverse
- (e) Range of inverse
- (f) Is the inverse a function?



9/21/16

HL1 Math - Santowski

3

Faster Five – Skills Preview

- For the following pairs of functions

- (a) Determine $f \circ g(x)$

(a) $f(x) = 3x - 6$ and $g(x) = \frac{1}{3}x + 2$

- (b) Determine $g \circ f(x)$

(b) $f(x) = \frac{1}{x+3}$ and $g(x) = \frac{1-3x}{x}$

- (c) Graph the original two functions in a square view window & make observations about the graph → then relate these observations back to the composition result

(c) $f(x) = 3 - (x+2)^2$ where $x \geq -2$ and $g(x) = \sqrt{3-x} - 2$

(d) $f(x) = e^{2x-1}$ and $g(x) = \frac{1}{2}(\ln(x)-1)$

9/21/16

HL1 Math - Santowski

4

Lesson Objectives

- Find the inverse of a function from numeric/tabular, graphic or algebraic data
- Compose a function with its inverse to develop the identity function
- Understand inverses as transformations

9/21/16

HL1 Math - Santowski

5

The **BIG** Picture

QUESTIONS??

- Are all functions invertible?
- Do function inverses “do the same thing” as our additive/multiplicative inverses?
- Why “invert” a function in the first place?

9/21/16

HL1 Math - Santowski

6

(A) Inverses – The Concept

- Let's back to our input \rightarrow output notion for functions.
- If functions are nothing more than input/output operators, then the concept of an inverse has us considering **how to go in reverse** \rightarrow going from the output back to the input

9/21/16

HL1 Math - Santowski

7

(A) Inverses – The Concept

- If the elements of the ordered pairs or mappings of a function are **reversed**, the resulting set of ordered pairs or mappings are referred to as the **INVERSE**.
- Since we are **REVERSING** the elements \rightarrow another point worth noting: the domain of the original function now becomes the range of the inverse; likewise, the range of the original becomes the domain of the inverse.

9/21/16


HL1 Math - Santowski

8

(A) Inverses – The Concept

Determine:

- Domain of $f(x)$
- Range of $f(x)$
- Mapping of inverse
- Domain of inverse
- Range of inverse
- Is the inverse a function?



9/21/16

HL1 Math - Santowski

9

(B) Notation of the Inverses

- If the inverse relation IS a function, then the notation used for these inverses is $f^{-1}(x)$.
- IMPORTANT NOTE:**
- $f^{-1}(x)$ does not mean $(f(x))^{-1}$ or $1/f(x)$.


9/21/16

HL1 Math - Santowski

10

(C) Examples

- Consider a graph of the following data:
 - Here is the graph of $f(x)$
- State domain and range of f
 - Evaluate $f(-2)$, $f(0)$,
 - Graph the inverse relation
 - Is the inverse a function?
 - HOW can we make the inverse a function?**
 - Evaluate $f^{-1}(1)$, $f^{-1}(-2)$
 - State the domain and range of $f^{-1}(x)$




9/21/16

HL1 Math - Santowski

11

(C) Examples

- Given the function of $y = f(x)$, determine
- the domain and range of the relation
 - the domain and range of the inverse relation
 - Is the inverse relation a function?
- Evaluate $f \circ f^{-1}(8)$
 - Evaluate $f(4)$
 - Solve $f(x) = \frac{1}{2}$
 - Evaluate $f^{-1}(1)$
 - Solve $f^{-1}(x) = 7$



9/21/16

HL1 Math - Santowski

12

(C) Examples

- Determine the equation for the inverse of the following functions. Draw both graphs and find the D and R of each.
- (1) $y = 4x - 9$
- (2) $y = 2x^2 + 4$ but
- (3) $y = 2 - \sqrt{x+3}$
- And some rational functions
- (5) $f(x) = \frac{2}{3x-9}$
- (6) $f(x) = \frac{2x-1}{3x-9}$
- (7) $y = |x|$

9/21/16

HL1 Math - Santowski

13

(C) Examples

- Given the function, determine $f(x) = 1 - \sqrt{x-2}$
- (a) the domain and range of the relation
- (b) the domain and range of the inverse relation
- (c) Determine the equation of the inverse relation
- (d) Is the inverse relation a function?
- (e) Evaluate $f^{-1}(5)$
- (f) Evaluate $f(6)$
- (g) Solve $f(x) = -2$
- (h) Evaluate $f^{-1}(-3)$
- (i) Solve $f^{-1}(x) = 11$

9/21/16

HL1 Math - Santowski

14

(D) Existence of Inverses

- A function, f , has an inverse, f^{-1} , if and only if f is one to one.
- So how do we "test" if a function is one to one?
- Let a be any real number, so if $x = a$, then $y = f(a)$
- Let b be any real number, so if $x = b$, then $y = f(b)$
- Recall that is $f(x)$ is one to one, then one output is produced by one input AND ALSO one input produces one output
- So, if our two outputs ($f(a)$ & $f(b)$) are equal AND if our two inputs are NOT the same, then our function is NOT one to one

9/21/16

HL1 Math - Santowski

15

(D) Existence of Inverses

- Let $f(x) = 2 + \sqrt{x}$
- So $f(a) = 2 + \sqrt{a}$
- And $f(b) = 2 + \sqrt{b}$
- If $f(a) = f(b)$
- Then $2 + \sqrt{a} = 2 + \sqrt{b}$
- So $\sqrt{a} = \sqrt{b}$
- Meaning that $a = b$
- so the only way to get the two same output values is to have the two input values to be identical, meaning you MUST be one to one.
- Let $f(x) = 2 + x^2$
- So $f(a) = 2 + a^2$
- And $f(b) = 2 + b^2$
- If $f(a) = f(b)$
- Then $2 + a^2 = 2 + b^2$
- So $a^2 = b^2$
- Now square root each side
- $\sqrt{a^2} = \sqrt{b^2}$
- So $|a| = |b|$
- Thus $a = \pm b$ or $b = \pm a$
- So now the SAME output value can be produced by different input values → NOT one to one!!

9/21/16

HL1 Math - Santowski

16

(D) Existence of Inverses

- Given $f(x) = \frac{x}{\sqrt{1+x^2}}$
- Show that $y = f(x)$ is a one to one function and hence find the equation of its inverse.

9/21/16

HL1 Math - Santowski

17

(D) Existence of Inverses

- Video resources for testing one to one:
- <https://youtu.be/9b8Xd77f-qk> from Krista King
- <https://youtu.be/TcSL2R360Aw> from Tom Killoran
- Reading resources for testing one to one
- <http://math.stackexchange.com/questions/101975/how-to-determine-if-a-function-is-one-to-one>
- http://www.math.toronto.edu/preparing-for-calculus/4_functions/we_3_one_to_one.html

9/22/16

HL1 Math - Santowski

18

(E) Composing with Inverses

- Let $f(x) = 2x - 7$.
- Determine the inverse of $y = f(x)$
- Graph both functions DESMOS
- Draw the line $y = x$. What do you observe? Why?
- What transformation are we considering in this scenario?
- Now compose as follows $f \circ f^{-1}(x)$ and $f^{-1} \circ f(x)$. What do you notice?

HL1 Math - Santowski

9/21/16
19

(E) Composing with Inverses

- Now let $f(x) = x^2 + 2$, **provided that** ???
- Determine the inverse of $y = f(x)$
- Graph both functions on DESMOS
- Draw the line $y = x$. What do you observe? Why?
- What transformation are we considering in this scenario?
- Now compose as follows $f \circ f^{-1}(x)$ and $f^{-1} \circ f(x)$. What do you notice?

9/21/16

HL1 Math - Santowski

20

(E) Composing with Inverses

- How can we use this observation?
- Determine the equation of the inverse of

$$f(x) = \frac{5}{x-3}$$

- Verify that your equation for the inverse IS correct (HINT: Composition)

9/21/16

HL1 Math - Santowski

21

Identity Functions & Inverses

Before we start our discussion of inverse functions, it is worthwhile looking back at a fundamental area of algebra – algebraic operations. The relevant algebraic properties for real numbers $a \in \mathbb{R}$, $b \in \mathbb{R}$ and $c \in \mathbb{R}$ are:

	Under addition	Under multiplication
Closure	$a + b \in \mathbb{R}$	$a \times b \in \mathbb{R}$
Commutativity	$a + b = b + a$	$a \times b = b \times a$
Associativity	$(a + b) + c = a + (b + c)$	$(a \times b) \times c = a \times (b \times c)$
Existence of the identity	$0 : a + 0 = 0 + a = a$	$1 : a \times 1 = 1 \times a = a$
Inverse element	$-a : a + (-a) = 0 = (-a) + a$	$\frac{1}{a} : \left(\frac{1}{a}\right) \times a = 1 = a \times \left(\frac{1}{a}\right)$

9/21/16

HL1 Math - Santowski

22

Identity Functions & Inverses

- Working with the Identity function & Inverses:

$$(f \circ I)(x) = f(x)$$

$$(I \circ f)(x) = f(x)$$

$$(f \circ f^{-1})(x) = \dots\dots$$

$$(f^{-1} \circ f)(x) = \dots\dots$$

9/21/16

HL1 Math - Santowski

23

(D) Exponential Functions

- Graph the exponential function $y = e^x + 1$ using DESMOS
- Now graph the inverse using DESMOS as well $\rightarrow x = e^y + 1$
- Explore the data tables of each function & the graphs & notice the connections
- Now graph the function $y = \ln(x - 1)$ and compare it to the graph of $x - 1 = e^y$
- Explain the point to the natural log function

9/21/16

HL1 Math - Santowski

24

(D) Exponential Functions

- Find the inverses of the following functions:

(a) $y = e^{2x}$

(b) $y = e^{x+2}$

(c) $y = 2e^x$

(d) $y = e^x + 2$

(e) $y = \ln(x+3) - 1$

(f) $y = 4\ln(2x)$