

Lesson 49 – Volumes of Revolution

Calculus - Santowski

Calculus - Santowski 4/22/2017

Lesson Objectives

- ▶ 1. Determine the volume of revolution of an object rotated about the x-axis
- ▶ 2. Determine by slicing (disk and washer method) or cylindrical shells to calculate volumes of solids
- ▶ 3. Apply volumes and average values to a real world problems

▶ 2 Calculus - Santowski 4/22/2017

Fast Five – Investigation

- ▶ 1. Determine the area of a circle if the radius is 6 cm.
- ▶ 2. Determine the area of a circle if its radius is defined by $y = 2x$ at the point where $x = 3$.
- ▶ 3. Draw the function $f(x) = 2x$ on the interval $[0,3]$. Estimate the area under $f(x)$ on $[0,3]$ using RRAM and 3 rectangles. Draw a diagram
- ▶ 4. Explain what happens when each of the 3 rectangles is completely rotated around the x-axis. Draw a diagram.
- ▶ 5. Explain what the idea of “volume of revolution” means

▶ 3 Calculus - Santowski 4/22/2017

(A) Volumes of Revolution

- ▶ Go to <http://archives.math.utk.edu/visual.calculus/5/volumes.5/index.html> and watch the animation showing the rotation of a graph about the x-axis and explaining how to determine the volume of the solid obtained in the animation above

- ▶ Explain the following formula:

$$V_x = \int_a^b A(x) dx = \int_a^b \pi \times (f(x))^2 dx = \pi \int_a^b (f(x))^2 dx$$

$$V_y = \int_a^b A(y) dy = \int_a^b \pi \times (f(y))^2 dy = \pi \int_a^b (f(y))^2 dy$$

▶ 4

Calculus - Santowski 4/22/2017

Example #1

- ▶ Go to <http://archives.math.utk.edu/visual.calculus/5/volumes.5/index.html> and consider the first example:
- ▶ Find the volume of the solid obtained by rotating the area bounded by the graph of $f(x) = x - x^2$ and the x-axis.

▶ 5

Calculus - Santowski 4/22/2017

Example #2

- ▶ Go to <http://archives.math.utk.edu/visual.calculus/5/volumes.5/index.html> and consider the second example:
- ▶ Find the volume of the solid obtained by rotating the area bounded by the graph of $f(x) = x^3 - x + 1$, $x = -1$, $x = 1$ and the x-axis.

▶ 6

Calculus - Santowski 4/22/2017

(D) Example 3

- ▶
- ▶ (b) Example 3: Area bounded by the graphs of $f(x) = x^3 - x + 1$, $x = -1$, $x = 1$ and the x-axis.

▶ ANS: $226\pi/105$

▶ <http://archives.math.utk.edu/visual.calculus/5/volumes.5/index.html>

▶

▶ 7

Calculus - Santowski 4/22/2017

(C) Example 4

- ▶ Example 4: Determine the volume of the solid obtained by rotating the region bounded by
- ▶ $f(x) = x^2 - 4x + 5$, $x = 1$, $x = 4$, and the x-axis about the x-axis.

▶ ANS: $78\pi/5$

▶ <http://tutorial.math.lamar.edu/Classes/Calcl/VolumeWithRings.aspx>

▶ 8

Calculus - Santowski 4/22/2017

(E) Example 5

- ▶ Determine the volume of the solid formed when $y = x^2$ is rotated around the y-axis between $y = 0$ and $y = 9$

▶ 9

Calculus - Santowski 4/22/2017

(E) Example 6

- ▶ Determine the volume of the solid formed when $y = x^2$ is rotated around the line $x = 1$ between $x = 1$ and $x = 3$

▶ 10

Calculus - Santowski 4/22/2017

(F) Volumes of Revolution – Rings & 2 Curves

- ▶ AREA of a RING → a region bounded by 2 curves
- ▶ $Area = \pi \times (\text{outer})^2 - \pi(\text{inner})^2 = \pi((\text{outer})^2 - (\text{inner})^2)$
- ▶ Formula to use: → see animation on <http://archives.math.utk.edu/visual.calculus/5/volume.s.5/index.html> (ring)

$$V_x = \int_a^b A(x) dx = \int_a^b \pi \times [(f(x))^2 - (g(x))^2] dx = \pi \int_a^b [(f(x))^2 - (g(x))^2] dx$$

$$V_y = \int_a^b A(y) dy = \int_a^b \pi \times [(f(y))^2 - (g(y))^2] dy = \pi \int_a^b [(f(y))^2 - (g(y))^2] dy$$

▶ 11

Calculus - Santowski 4/22/2017

(G) Example 1

- ▶ (c) Example 1: Determine the volume of the solid obtained by rotating the portion of the region bounded by the following 2 curves that lies in the first quadrant about the x-axis.

$$y = \sqrt[3]{x} \quad \text{and} \quad y = \frac{x}{4}$$

- ▶ ANS: $128\pi/15$

- ▶ <http://tutorial.math.lamar.edu/Classes/Calci/VolumeWithRings.aspx>

▶ 12

Calculus - Santowski 4/22/2017

(H) Example 2

- ▶ Find the volume of the solid obtained by rotating the area bounded by $f(x) = x^2$ and $g(x) = x$ about the line $y = 2$.
- ▶ ANS: $8\pi/15$
- ▶ <http://archives.math.utk.edu/visual.calculus/5/volumes.5/index.html>

▶ 13

Calculus - Santowski 4/22/2017

(I) Example 3

- ▶ Determine the volume of the solid obtained by rotating the region bounded by the functions $y = x$ and $y = x^2 - 2x$ about the line $y = 4$.
- ▶ ANS: $153\pi/5$
- ▶ <http://tutorial.math.lamar.edu/Classes/Calcl/VolumeWithRings.aspx>

▶ 14

Calculus - Santowski 4/22/2017

(J) Example 4

- ▶ Determine the volume of the solid obtained by rotating the region bounded by $y = x - 1$ and $y = 2\sqrt{x-1}$ and about the line $x = -1$.
- ▶ ANS: $96\pi/5$
- ▶ <http://tutorial.math.lamar.edu/Classes/Calcl/VolumeWithRings.aspx>

▶ 15

Calculus - Santowski 4/22/2017
