

# Lesson 71 – Introduction to Complex Numbers

HL2 MATH - SANTOWSKI

---

---

---

---

---

---

---

## Lesson Objectives

- (1) Introduce the idea of imaginary and complex numbers
- (2) Practice operations with complex numbers
- (3) Use complex numbers to solve polynomials
- (4) geometric representation of complex numbers

---

---

---

---

---

---

---

To see a complex number we have to  
first see where it shows up

Solve both of these

$$x^2 - 81 = 0$$

$$x^2 = 81$$

$$x = \pm 9$$

$$x^2 + 81 = 0$$

$$x^2 = -81$$




---

---

---

---

---

---

---

Um, no solution????

$x = \pm\sqrt{-81}$  does not have a real answer.

It has an "imaginary" answer.

To define a complex number we have to create a new variable.

This new variable is "i"

---

---

---

---

---

---

---

## Imaginary Unit

Until now, you have always been told that you can't take the square root of a negative number. If you use imaginary units, you can!

The imaginary unit is  $i$  where  $i = \sqrt{-1}$

It is used to write the square root of a negative number.

---

---

---

---

---

---

---

## Property of the square root of negative numbers

If  $r$  is a positive real number, then  $\sqrt{-r} = i\sqrt{r}$

Examples:

$$\sqrt{-3} = i\sqrt{3}$$

$$\sqrt{-4} = i\sqrt{4} = 2i$$

---

---

---

---

---

---

---

Definition:  $i = \sqrt{-1}$

Note:  $i$  is the representation for  $\sqrt{-1}$ , not a simplification of  $\sqrt{-1}$

So, following this definition:  $i^2 = -1$

So what is  $i^3$  and  $i^4$ ?

---

---

---

---

---

---

---

And it cycles....

$i = \sqrt{-1}$	$i^5 = i^4 \cdot i = i$	$i^9 = i^8 \cdot i = i$
$i^2 = -1$	$i^6 = i^4 \cdot i^2 = -1$	$i^{10} = i^8 \cdot i^2 = -1$
$i^3 = -i$	$i^7 = i^4 \cdot i^3 = -i$	$i^{11} = i^8 \cdot i^3 = -i$
$i^4 = 1$	$i^8 = i^4 \cdot i^4 = 1$	$i^{12} = i^8 \cdot i^4 = 1$

Do you see a pattern yet?

---

---

---

---

---

---

---

What is that pattern?

We are looking at the remainder when the power is divided by 4.

Why?

Every  $i^4$  doesn't matter. It is what remains after all of the  $i^4$  are taken out.

Try it with  $i^{92233}$

---

---

---

---

---

---

---

## Integral powers of i(iota)

$$i^0 = 1 \text{ (as usual)}$$

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = i^2 \cdot i = -i$$

$$i^4 = i^3 \cdot i = -i \cdot i = 1$$

$$i^{-1} = \frac{1}{i} = \frac{1}{i} \cdot \frac{i}{i} = -i$$

$$i^{-2} = \frac{1}{i^2} = -1$$

$$i^{-3} = \frac{1}{i^3} = \frac{1}{-i} = i$$

$$i^{-4} = \frac{1}{i^4} = 1$$

Evaluate:

$$\left( i^{17} - \left( \frac{2}{i} \right)^3 \right)$$

## Integral powers of i(iota)

$$i^0 = 1 \text{ (as usual)}$$

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = i^2 \cdot i = -i$$

$$i^4 = i^3 \cdot i = -i \cdot i = 1$$

$$i^{-1} = \frac{1}{i} = \frac{1}{i} \cdot \frac{i}{i} = -i$$

$$i^{-2} = \frac{1}{i^2} = -1$$

$$i^{-3} = \frac{1}{i^3} = \frac{1}{-i} = i$$

$$i^{-4} = \frac{1}{i^4} = 1$$

Evaluate:

$$\left( i^{17} - \left( \frac{2}{i} \right)^3 \right)$$

Solution

$$\left( i^{16} \cdot i - \frac{8}{i^3} \right) = \left( i + \frac{8}{i} \right) = (i - 8i)$$

Ans: 343i

## Illustrative Problem

If p, q, r, s are four consecutive integers, then  $i^p + i^q + i^r + i^s =$

- a) 1      b) 2  
c) 4      d) None of these

### Illustrative Problem

If  $p, q, r, s$  are four consecutive integers, then  $i^p + i^q + i^r + i^s =$

- a) 1                  b) 2  
c) 4                  d) None of these

Solution: Note  $q = p + 1, r = p + 2, s = p + 3$

Given expression =  $i^p(1 + i + i^2 + i^3)$

=  $i^p(1 + i - 1 - i) = 0$                   Remember this.

---

---

---

---

---

---

---

---

### Illustrative Problem

If  $u_{n+1} = i u_n + 1$ , where  $u_1 = i + 1$ , then  $u_{27}$  is

- a)  $i$                   b) 1  
c)  $i + 1$               d) 0

---

---

---

---

---

---

---

---

### Illustrative Problem

If  $u_{n+1} = i u_n + 1$ , where  $u_1 = i + 1$ , then  $u_{27}$  is

- a)  $i$                   b) 1  
c)  $i + 1$               d) 0

Solution:  $u_2 = i u_1 + 1 = i(i+1) + 1 = i^2 + i + 1$

$u_3 = i u_2 + 1 = i(i^2 + i + 1) + 1 = i^3 + i^2 + i + 1$

Hence  $u_n = i^n + i^{n-1} + \dots + i + 1$                   Note by previous question:

$u_{27} = i^{27} + i^{26} + \dots + i + 1 = \frac{i^{28} - 1}{i - 1} = 0$                    $u_{27} = 0$

---

---

---

---

---

---

---

---

## Hints to deal with $i$

1. Find all " $i$ 's at the beginning of a problem.
2. Treat all " $i$ 's like variables, with all rules of exponents holding.
3. Reduce the power of  $i$  at the end by the rules we just learned..

---

---

---

---

---

---

---

## Examples

1.  $\sqrt{-36} \cdot \sqrt{-81}$

2.  $\sqrt{-36} + \sqrt{-81}$

---

---

---

---

---

---

---

## COMPLEX NUMBERS

But what is  $1 + 3i$

The two types of number ( $1$  and  $3i$ ) cannot be "mixed".

Numbers of the form  $k \times i$ ,  $k \in \mathbb{R}$  are called **imaginary numbers** (or "pure imaginary")

Numbers like  $1$ ,  $2$ ,  $-3.8$  that we used before are called **real numbers**.

When we combine them together in a sum we have **complex numbers**.

---

---

---

---

---


---

---

OK, so what is a complex number?

A complex number has two parts – a real part and an imaginary part.

A complex number comes in the form  $a + bi$




---

---

---

---

---

---

---

#### COMPLEX NUMBERS

To summarize,

$$z = a + bi$$

- $a$  and  $b$  are real numbers
- $a$  is the "**real part**" of  $z$ ;  $\text{Re}(z)$
- $b$  is the "**imaginary part**" of  $z$ ;  $\text{Im}(z)$
- The sum of the two parts is called a "**complex number**"

---

---

---

---

---

---

---

And just so you know...

All real numbers are complex  $\rightarrow 3 = 3 + 0i$

All imaginary numbers are complex  $\rightarrow 7i = 0 + 7i$

Again, treat the  $i$  as a variable and you will have no problems.

---

---

---

---

---

---

---

## COMPLEX NUMBERS

Adding and subtracting complex numbers:

$$z_1 = (2 + 3i)$$

$$z_2 = (4 - 9i)$$

---

---

---

---

---

---

---

## COMPLEX NUMBERS

Adding and subtracting complex numbers:

$$z_1 = (2 + 3i)$$

$$z_2 = (4 - 9i) \quad z_1 + z_2 = 6 - 6i$$

$$(a + bi) \pm (c + di) \equiv (a \pm c) + (b \pm d)i$$

For addition and subtraction the real and imaginary parts are kept separate.

---

---

---

---

---

---

---

Adding and Subtracting  
(add or subtract the real parts, then add or subtract the imaginary parts)

Ex:  $(-1 + 2i) + (3 + 3i)$

Ex:  $2i - (3 + i) + (2 - 3i)$

Ex:  $(2 - 3i) - (3 - 7i)$

---

---

---

---

---

---

---



Adding and Subtracting

(add or subtract the real parts, then add or subtract the imaginary parts)

$$\begin{array}{ll} \text{Ex: } (-1+2i) + (3+3i) & \text{Ex: } 2i - (3+i) + (2-3i) \\ = (-1+3) + (2i+3i) & = (-3+2) + (2i-i-3i) \\ = 2+5i & = -1-2i \\ \\ \text{Ex: } (2-3i) - (3-7i) & \\ = (2-3) + (-3i-(-7i)) & \\ = -1+4i & \end{array}$$

---

---

---

---

---

---

---

---

## COMPLEX NUMBERS

Multiplying and dividing complex numbers:

$$\begin{aligned} z_1 &= (2+3i) \\ z_2 &= (4-9i) \end{aligned}$$

---

---

---

---

---

---

---

---

## COMPLEX NUMBERS

Multiplying and dividing complex numbers:

$$\begin{aligned} z_1 &= (2+3i) & z_1 z_2 &= (2+3i) \times (4-9i) \\ z_2 &= (4-9i) & &= 2 \times 4 + (2 \times -9i) + (3i \times 4) + (3i \times -9i) \\ & & &= 8 - 18i + 12i + (-27 \times i^2) \\ & & &= 35 - 6i \end{aligned}$$

$$(a+bi) \times (c+di) \equiv (ac-bd) + (bc+ad)i$$

Notice how, for multiplication, the real and imaginary parts "mix" through the formula  $i^2 = -1$ .

---

---

---

---

---

---

---

---

### Multiplying

Ex:  $-i(3+i)$

Ex:  $(2+3i)(-6-2i)$

---

---

---

---

---

---

---

Multiplying → Treat the i's like variables, then change any that are not to the first power

Ex:  $-i(3+i)$   
 $= -3i - i^2$   
 $= -3i - (-1)$   
 $= 1 - 3i$

Ex:  $(2+3i)(-6-2i)$   
 $= -12 - 4i - 18i - 6i^2$   
 $= -12 - 22i - 6(-1)$   
 $= -12 - 22i + 6$   
 $= -6 - 22i$

---

---

---

---

---

---

---

### COMPLEX CONJUGATES

What are the solutions to  $x^2 - 6x + 21 = 0$  ?

$$3 \pm 2\sqrt{3}i$$

If we write  $z = 3 + 2\sqrt{3}i$

Then the complex conjugate is written as  $z^* = 3 - 2\sqrt{3}i$

Calculate the following:

$$z + z^*$$

$$z - z^*$$

$$zz^*$$

---

---

---

---

---

---

---

## COMPLEX CONJUGATES

What are the solutions to  $x^2 - 6x + 21 = 0$  ?

$$3 \pm 2\sqrt{3}i$$

If we write  $z = 3 + 2\sqrt{3}i$

\* means conjugate

Then the complex conjugate is written as  $z^* = 3 - 2\sqrt{3}i$

Calculate the following:

$$z + z^* = 6 = 2\operatorname{Re}(z)$$

$$z - z^* = 4\sqrt{3}i = 2\operatorname{Im}(z)$$

$$zz^* = 3^2 + (2\sqrt{3})^2 = 21 = |z|^2$$

## COMPLEX NUMBERS

Dividing complex numbers:

$$\frac{z_1}{z_2} = \frac{(2+3i)}{(4-9i)}$$

$$\frac{z_1}{z_2} =$$

## COMPLEX NUMBERS

Dividing complex numbers:

$$\frac{z_1}{z_2} = \frac{(2+3i)}{(4-9i)}$$

$$\frac{z_1}{z_2} =$$

$$\begin{aligned} &= \frac{(2+3i)}{(4-9i)} \times \frac{(4+9i)}{(4+9i)} \\ &= \frac{8+18i+12i+(27 \times i^2)}{4 \times 4 + 36i - 36i + (-9 \times 9 \times i^2)} \\ &= \frac{-19+30i}{97} = -\frac{19}{97} + \frac{30}{97}i \end{aligned}$$

Remember this trick!!

$$\text{Ex: } \frac{3+11i}{-1-2i}$$

---

---

---

---

---

---

---

$$\begin{aligned} \text{Ex: } \frac{3+11i}{-1-2i} \cdot \frac{-1+2i}{-1+2i} &= \frac{-25-5i}{5} \\ &= \frac{(3+11i)(-1+2i)}{(-1-2i)(-1+2i)} = \frac{-25-5i}{5} \\ &= \frac{-3+6i-11i+22i^2}{1-2i+2i-4i^2} = \frac{-3-5i-22}{1-4(-1)} \\ &= \frac{-3-5i+22(-1)}{1-4(-1)} = \frac{-3-5i-22}{1+4} \\ &= \frac{-25-5i}{5} \end{aligned}$$

$$= \frac{-25-5i}{5} = -5-i$$

---

---

---

---

---

---

---

### More Practice

5.  $6i^{-5}$

6.  $\frac{6-i}{4} + \frac{4+2i}{3+i}$

---

---

---

---

---

---

---

## Absolute Value of a Complex Number

The distance the complex number is from the origin on the complex plane.

If you have a complex number  $(a + bi)$

the absolute value can be found using:  $\sqrt{a^2 + b^2}$

---

---

---

---

---

---

---

---

## Examples

1.  $|-2 + 5i|$       2.  $|-6i|$

---

---

---

---

---

---

---

---

## Examples

1.  $|-2 + 5i|$   
 $= \sqrt{(-2)^2 + (5)^2}$   
 $= \sqrt{4 + 25}$   
 $= \sqrt{29}$
2.  $|-6i|$   
 $= \sqrt{(0)^2 + (-6)^2}$   
 $= \sqrt{0 + 36}$   
 $= \sqrt{36}$   
 $= 6$

Which of these 2 complex numbers is closest to the origin?

$-2 + 5i$

---

---

---

---

---

---

---

---

### Complex Conjugates Theorem

Roots/Zeros that are not *Real* are *Complex* with an *Imaginary* component. Complex roots with Imaginary components always exist in Conjugate Pairs.

If  $a + bi$  ( $b \neq 0$ ) is a zero of a polynomial function, then its Conjugate,  $a - bi$ , is also a zero of the function.

---

---

---

---

---

---

---

---

### Find Roots/Zeros of a Polynomial

If the known root is *imaginary*, we can use the *Complex Conjugates Theorem*.

Ex: Find all the roots of  $f(x) = x^3 - 5x^2 - 7x + 51$

If one root is  $4 - i$ .

Because of the Complex Conjugate Theorem, we know that *another* root must be  $4 + i$ .

Can the third root also be imaginary?

---

---

---

---

---

---

---

---

### Example (con't)

Ex: Find all the roots of  $f(x) = x^3 - 5x^2 - 7x + 51$

If one root is  $4 - i$ .

If one root is  $4 - i$ , then one factor is  $[x - (4 - i)]$ , and

Another root is  $4 + i$ , & another factor is  $[x - (4 + i)]$ .

Multiply these factors:

$$\begin{aligned} [x - (4 - i)] [x - (4 + i)] &= x^2 - x(4 + i) - x(4 - i) + (4 - i)(4 + i) \\ &= x^2 - 4x - xi - 4x + xi + 16 - i^2 \\ &= x^2 - 8x + 16 - (-1) \\ &= x^2 - 8x + 17 \end{aligned}$$

---

---

---

---

---

---

---

---

### Example (con't)

Ex: Find all the roots of  $f(x) = x^3 - 5x^2 - 7x + 51$

If one root is  $4 - i$ .

If the product of the two non-real factors is  $x^2 - 8x + 17$   
then the third factor (that gives us the neg. real root) is  
the quotient of  $P(x)$  divided by  $x^2 - 8x + 17$ :

$$\begin{array}{r} x^2 - 8x + 17 \overline{) x^3 - 5x^2 - 7x + 51} \\ \underline{x^3 - 8x^2 + 17x - 51} \phantom{0} \\ 3x^2 - 24x + 102 \phantom{0} \\ \underline{3x^2 - 24x + 51} \\ 51 \phantom{0} \\ \underline{51} \\ 0 \end{array}$$

The third root  
is  $x = -3$

Now write a polynomial function of least degree that has **real coefficients**, a **leading coeff. of 1** and **1, -2+i, -2-i** as zeros.

Now write a polynomial function of least degree that has **real coefficients**, a **leading coeff. of 1** and **1, -2+i, -2-i** as zeros.

$$f(x) = (x-1)(x-(-2+i))(x-(-2-i))$$

$$f(x) = (x-1)(x+2-i)(x+2+i)$$

$$f(x) = (x-1)((x+2) - i)((x+2)+i)$$

$$f(x) = (x-1)((x+2)^2 - i^2)$$

Foil

$$f(x) = (x-1)(x^2 + 4x + 4 - (-1))$$

Take care of  $i^2$

$$f(x) = (x-1)(x^2 + 4x + 4 + 1)$$

$$f(x) = (x-1)(x^2 + 4x + 5)$$

Multiply

$$f(x) = x^3 + 4x^2 + 5x - x^2 - 4x - 5$$

$$f(x) = x^3 + 3x^2 + x - 5$$

Now write a polynomial function of least degree that has **real coefficients**, a **leading coeff. of 1** and **4, 4, 2+i** as zeros.

---

---

---

---

---

---

---

---

Now write a polynomial function of least degree that has **real coefficients**, a **leading coeff. of 1** and **4, 4, 2+i** as zeros.

Note:  $2+i$  means  $2-i$  is also a zero

$$F(x) = (x-4)(x-4)(x-(2+i))(x-(2-i))$$

$$F(x) = (x-4)(x-4)(x-2-i)(x-2+i)$$

$$F(x) = (x^2 - 8x + 16)[(x-2) - i][(x-2) + i]$$

$$F(x) = (x^2 - 8x + 16)[(x-2)^2 - i^2]$$

$$F(x) = (x^2 - 8x + 16)(x^2 - 4x + 4 - (-1))$$

$$F(x) = (x^2 - 8x + 16)(x^2 - 4x + 5)$$

$$F(x) = x^4 - 4x^3 + 5x^2 - 8x^3 + 32x^2 - 40x + 16x^2 - 64x + 80$$

$$F(x) = x^4 - 12x^3 + 53x^2 - 104x + 80$$

---

---

---

---

---

---

---

---

## Further Examples

**EXAMPLES:** Find a polynomial with the given zeros

-1, -1, 3i, -3i

2, 4 + i, 4 - i

---

---

---

---

---

---

---

---



**EXAMPLE:** Solving a Polynomial Equation

Solve:  $x^4 - 6x^2 - 8x + 24 = 0$ .

---

---

---

---

---

---

---

**EXAMPLE:** Solving a Polynomial Equation

Solve:  $x^4 - 6x^2 - 8x + 24 = 0$ .

**Solution** Now we can solve the original equation as follows.

$$x^4 - 6x^2 + 8x + 24 = 0 \text{ This is the given equation.}$$

$$(x-2)(x-2)(x^2+4x+6) = 0 \quad \text{This was obtained from the second synthetic division.}$$

$$x-2=0 \text{ or } x-2=0 \text{ or } x^2+4x+6=0 \text{ Set each factor equal to zero.}$$

$$x=2 \quad x=2 \quad x^2+4x+6=0 \text{ Solve.}$$

---

---

---

---

---

---

---

**EXAMPLE:** Solving a Polynomial Equation

Solve:  $x^4 - 6x^2 - 8x + 24 = 0$ .

**Solution** We can use the quadratic formula to solve  $x^2 + 4x + 6 = 0$ .

$$\frac{-4 \pm \sqrt{4^2 - 4(1)(6)}}{2(1)}$$

$$\frac{-4 \pm \sqrt{16 - 24}}{2}$$

$$\frac{-4 \pm \sqrt{-8}}{2}$$

$$\frac{-4 \pm 2\sqrt{2}i}{2}$$

$$-2 \pm \sqrt{2}i$$

We use the quadratic formula because  $x^2 + 4x + 6 = 0$  cannot be factored.Let  $a=1$ ,  $b=4$ , and  $c=6$ .

Multiply and subtract under the radical.

$$\sqrt{4^2 - 4(1)(6)}$$

Simplify.

The solution set of the original equation is  $\{2, -2 \pm i, -2 \pm i\sqrt{2}\}$ .

---

---

---

---

---

---

---

### FIND ALL THE ZEROS

$$f(x) = x^4 - 3x^3 + 6x^2 + 2x - 60$$

(Given that  $1 + 3i$  is a zero of  $f$ )

$$f(x) = x^3 - 7x^2 - x + 87$$

(Given that  $5 + 2i$  is a zero of  $f$ )

---

---

---

---

---

---

---

### More Finding of Zeros

$$f(x) = x^5 + x^3 + 2x^2 - 12x + 8$$

$$f(x) = 3x^3 - 4x^2 + 8x + 8$$

---

---

---

---

---

---

---

Find the zeros of

Hint: 4 is a zero

$$f(x) = x^3 - 11x - 20$$

---

---

---


---

---

---

---

Find the zeros of  $f(x) = x^3 - 11x - 20$   
 Hint: 4 is a zero



$$\begin{array}{r|rrrrr} 4 & 1 & 0 & -11 & -20 & \\ & & 4 & 16 & 20 & \\ \hline & 1 & 4 & 5 & 0 & \end{array}$$

$$(x-4)(x^2+4x+5)=0$$

$$\frac{-4 \pm \sqrt{16-4(1)(5)}}{2}$$

$$\frac{-4 \pm \sqrt{-4}}{2}$$

$$-2 \pm i$$

$$x = 4, -2+i, -2-i$$


---

---

---

---

---

---

---

---

No Calculator  
 Given 2 is a zero of  $f(x) = x^3 - 6x^2 + 13x - 10$ ,  
 find ALL the zeros of the function.

---

---

---

---

---

---

---

---

No Calculator  
 Given 2 is a zero of  $f(x) = x^3 - 6x^2 + 13x - 10$ ,  
 find ALL the zeros of the function.

$$\begin{array}{r|rrrrr} 2 & 1 & -6 & 13 & -10 & \\ & & 2 & -8 & 10 & \\ \hline & 1 & -4 & 5 & 0 & \end{array}$$

$$(x-2)(x^2-4x+5)=0$$

$$\frac{4 \pm \sqrt{16-4(1)(5)}}{2}$$

$$\frac{4 \pm \sqrt{-4}}{2}$$

$$2 \pm i$$

$$x = 2, 2+i, 2-i$$


---

---

---

---

---

---

---

---

No Calculator

Given  $-3$  is a zero of  $f(x) = x^3 + 3x^2 + x + 3$ ,  
find ALL the zeros of the function.

---

---

---

---

---

---

---

No Calculator

Given  $-3$  is a zero of  $f(x) = x^3 + 3x^2 + x + 3$ ,  
find ALL the zeros of the function.

$$\begin{array}{r|rrrr} -3 & 1 & 3 & 1 & 3 \\ & & -3 & 0 & -3 \\ \hline & 1 & 0 & 1 & 0 \end{array}$$

$$(x+3)(x^2+1) = 0$$

$$x^2 = -1$$

$$x = i, -i$$

$$x = -3, i, -i$$

---

---

---

---

---

---

---