




LESSON 85 – DISTANCES IN 3 SPACE

HL2 - Santowski

Distance Between a Point and a Line

In the picture below, Q is a point not on the line, P is a point on the line, \vec{u} is a direction vector for the line and θ is the angle between \vec{u} and \vec{PQ} .



Obviously, $\sin \theta = \frac{D}{\|\vec{PQ}\|}$ or $D = \|\vec{PQ}\| \sin \theta$



We know from Section 7.4 on cross products that

$$\|\vec{u} \times \vec{v}\| = \|\vec{u}\| \|\vec{v}\| \sin \theta, \text{ where } \theta \text{ is the angle between } \vec{u} \text{ and } \vec{v}.$$

Thus, $\|\vec{PQ} \times \vec{u}\| = \|\vec{PQ}\| \|\vec{u}\| \sin \theta$

or dividing both sides by $\|\vec{u}\|$

$$\frac{\|\vec{PQ} \times \vec{u}\|}{\|\vec{u}\|} = \|\vec{PQ}\| \sin \theta$$

So if, $D = \|\vec{PQ}\| \sin \theta$ then from above, $D = \frac{\|\vec{PQ} \times \vec{u}\|}{\|\vec{u}\|}$.



Distance Between a Point and a Line

The distance, D , between a line and a point Q not on the line is given by

$$D = \frac{\|\vec{PQ} \times \vec{u}\|}{\|\vec{u}\|}$$

where \vec{u} is the direction vector of the line and P is a point on the line.

Example 1: Find the distance between the point $Q(1, 3, -2)$ and the line given by the parametric equations:

$$x = 2 + t, \quad y = -1 - t \quad \text{and} \quad z = 3 + 2t$$

Solution: From the parametric equations we know the direction vector, \vec{u} is $\langle 1, -1, 2 \rangle$ and if we let $t = 0$, a point P on the line is $P(2, -1, 3)$.

Thus $\vec{PQ} = \langle 2-1, -1-3, 3-(-2) \rangle = \langle 1, -4, 5 \rangle$

Find the cross product:

$$\vec{PQ} \times \vec{u} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & -4 & 5 \\ 1 & -1 & 2 \end{vmatrix} = -3\vec{i} + 3\vec{j} + 3\vec{k}$$

Using the distance formula:

$$D = \frac{\|\vec{PQ} \times \vec{u}\|}{\|\vec{u}\|} = \frac{\sqrt{(-3)^2 + 3^2 + 3^2}}{\sqrt{1^2 + (-1)^2 + 2^2}} = \frac{\sqrt{27}}{\sqrt{6}} = \sqrt{\frac{9}{2}} \approx 2.12$$

DISTANCES BETWEEN LINES

o Example #2:

o (a) Show that the lines are skew.

$$L1: x = 1 + t \quad y = -2 + 3t \quad z = 4 - t$$

$$L2: x = 2s \quad y = 3 + s \quad z = -3 + 4s$$

o (b) Find the distance between them.

DISTANCES BETWEEN LINES

- Since the two lines L_1 and L_2 are skew, they can be viewed as lying on two parallel planes P_1 and P_2 .
 - The distance between L_1 and L_2 is the same as the distance between P_1 and P_2 .



DISTANCES BETWEEN LINES

- The common normal vector to both planes must be orthogonal to both
- $v_1 = \langle 1, 3, -1 \rangle$ (direction of L_1)
- $v_2 = \langle 2, 1, 4 \rangle$ (direction of L_2)



DISTANCES BETWEEN LINES

- If we put $s = 0$ in the equations of L_2 , we get the point $(0, 3, -3)$ on L_2 .
 - So, an equation for P_2 is:
$$13(x - 0) - 6(y - 3) - 5(z + 3) = 0$$
or $13x - 6y - 5z + 3 = 0$



DISTANCES BETWEEN LINES

- If we now set $t = 0$ in the equations for L_1 , we get the point $(1, -2, 4)$ on P_1 .



DISTANCES BETWEEN LINES

- So, the distance between L_1 and L_2 is the same as the distance from $(1, -2, 4)$ to $13x - 6y - 5z + 3 = 0$.


$$d = \frac{8}{\sqrt{230}} = 0.53$$



Distance Between a Point and a Plane

Let P be a point in the plane and let Q be a point not in the plane. We are interested in finding the distance from the point Q to the plane that contains the point P .

We can find the distance between the point, Q , and the plane by projecting the vector from P to Q onto the normal to the plane and then finding its magnitude or length.



Thus the distance from Q to the plane is the length or the magnitude of the projection of the vector PQ onto the normal.



Distance Between a Point and a Plane

If the distance from Q to the plane is the length or the magnitude of the projection of the vector PQ onto the normal, we can write that mathematically:

$$\text{Distance from Q to the plane} = \|\text{proj}_{\vec{n}} \vec{PQ}\|$$

Now, recall from section 7.3, $\text{proj}_{\vec{n}} \vec{PQ} = \left(\frac{\vec{PQ} \cdot \vec{n}}{\|\vec{n}\|^2} \right) \cdot \vec{n}$

So taking the magnitude of this vector, we get:

$$\|\text{proj}_{\vec{n}} \vec{PQ}\| = \left\| \left(\frac{\vec{PQ} \cdot \vec{n}}{\|\vec{n}\|^2} \right) \cdot \vec{n} \right\| = \frac{|\vec{PQ} \cdot \vec{n}|}{\|\vec{n}\|^2} \cdot \|\vec{n}\| = \frac{|\vec{PQ} \cdot \vec{n}|}{\|\vec{n}\|}$$



Distance Between a Point and a Plane

The distance from a plane containing the point P to a point Q not in the plane is

$$D = \|\text{proj}_{\vec{n}} \vec{PQ}\| = \frac{|\vec{PQ} \cdot \vec{n}|}{\|\vec{n}\|}$$

where n is a normal to the plane.



Example 1: Find the distance between the point Q (3, 1, -5) to the plane $4x + 2y - z = 8$.

Solution: We know the normal to the plane is $\langle 4, 2, -1 \rangle$ from the general form of a plane. We can find a point in the plane simply by letting x and y equal 0 and solving for z: P (0, 0, -8) is a point in the plane.

Thus the vector, $\vec{PQ} = \langle 3-0, 1-0, -5-(-8) \rangle = \langle 3, 1, 3 \rangle$

Now that we have the vector \vec{PQ} and the normal, we simply use the formula for the distance between a point and a plane.

$$D = \|\text{proj}_{\vec{n}} \vec{PQ}\| = \frac{|\vec{PQ} \cdot \vec{n}|}{\|\vec{n}\|} = \frac{|\langle 3, 1, 3 \rangle \cdot \langle 4, 2, -1 \rangle|}{\sqrt{4^2 + 2^2 + (-1)^2}}$$

$$D = \frac{|12 + 2 - 3|}{\sqrt{16 + 4 + 1}} = \frac{11}{\sqrt{21}} \approx 2.4$$



Example 2: Find the distance between the point (1, 2, 3) and plane $2x - y + z = 4$

Solution:



DISTANCES BETWEEN POINTS & PLANES

- Find the distance between the parallel planes

$$10x + 2y - 2z = 5 \text{ and } 5x + y - z = 1$$



DISTANCES BETWEEN POINTS & PLANES

- First, we note that the planes are parallel because their normal vectors are parallel.

i.e. $\langle 10, 2, -2 \rangle$ and $\langle 5, 1, -1 \rangle$



DISTANCES BETWEEN POINTS & PLANES

- To find the distance D between the planes, we choose any point on one plane and calculate its distance to the other plane.
 - In particular, if we put $y = z = 0$ in the equation of the first plane, we get $10x = 5$.
 - So, $(\frac{1}{2}, 0, 0)$ is a point in this plane.



DISTANCES BETWEEN POINTS & PLANES

So, the distance between the planes is $\frac{\sqrt{3}}{6}$