



## Chapter Self-Test, page 64

1. Let  $x$  represent the number of 500 g cartons, and let  $y$  represent the number of 750 g cartons;  $0.5x + 0.75y = 887.5$

Possible Combinations  
of Raisin Bags




2. a) Let  $V$  represent the volume remaining, and let  $t$  represent the time after 8:30 a.m.;  $V = 1500 - 4t$   
 b) Answers may vary, e.g., about 10:30 a.m.  
 c) 10:35 a.m.
3. a)  $(-1.5, 2.5)$     b)  $(\frac{-24}{7}, \frac{1}{7})$     c)  $(2, -3)$
4. about 13.33 g of 85% gold, about 6.67 g of 70% gold
5. Answers may vary, e.g., at the point  $(x, y)$ , which represents a solution to a linear system, both sides of an equation in the system must be equal. Therefore, adding or subtracting these equations is the same as adding or subtracting constants to both sides of an equation: the solution will remain the same.
6. a)  $4x + 2y = 1, 2x + 6y = -17; x = 2, y = -3.5$   
 b)  $4x + 2y = 1, 2x + 6y = -17, x = 2,$   
 $y = -3.5, 3x + 4y = -8, x - 2y = 9$




7. 6 km  
 8. \$1500 in a savings account, \$2700 in bonds  
 9. Answers may vary, e.g., adding the first equation to 3 times the second equation and then simplifying gives  $15 = 24$ , which is not true.

5. a)  $-2$     c) 8    e) 3 or  $-3$   
 b)  $-1$     d) 6 or  $-6$     f) 8 or  $-8$
6. a)  $(1, 7)$     b)  $(1, \frac{3}{2})$
7. a) 6    b)  $\frac{1}{4}$     c) 0.7
8. a) about  $36.2 \text{ cm}^2$   
 b) about  $57.0 \text{ cm}^2$ , about  $41.7 \text{ cm}$
- 9.




## Lesson 2.1, page 78

1. a)  $(3, 5)$     b)  $(-0.5, 3.5)$   
 2. a)  $(2, 3)$     b)  $(0.5, 1)$     c)  $(4, -2)$   
 3. a)



- b)  $(32, 19)$
4. a)  $(2, 5)$     c)  $(2, -2)$     e)  $(-1, -1)$   
 b)  $(0.5, 3.5)$     d)  $(-0.5, 0.5)$     f)  $(0.25, 0.75)$
5.  $(0.75, -1)$
6.  $(5, 3)$ ; from  $P$  to  $M$ , run = 4 and rise = 2; the run and rise will be the same from  $M$  to  $Q$ , so  $Q$  has coordinates  $(1 + 4, 1 + 2)$
7. a), b)  $y = -\frac{1}{2}x - 1$



8. Answers may vary, e.g.,  $(-4, 4)$  and  $(2, 2)$  based on the assumption that the centre is at  $O$ , or  $(5, 1)$  and  $(-1, 3)$  based on the assumption that the centre is at  $R$

## Chapter 2

### Getting Started, page 68


1. a) viii    c) ii    e) iv    g) i  
 b) vii    d) v    f) vi    h) iii
2. a) 13 m  
 b) about 192.3 mm
3. a)  $y = \frac{1}{3}x + \frac{14}{3}$   
 b)  $y = -4x - 6$   
 c)  $y = -5x + 17$
4. a)  $-\frac{3}{2}$     c)  $\frac{35}{3}x$     e)  $\frac{23}{20}$   
 b)  $-\frac{3}{56}$     d)  $\frac{3}{8}y$     f)  $-1.4375$






- b) i) 7 units    ii) (7, 0), (-7, 0), (0, 7), (0, -7)

iii)




- c) i) 0.2 units    ii) (0.2, 0), (-0.2, 0), (0, 0.2), (0, -0.2)

iii)




- d) i) 13 units    ii) (13, 0), (-13, 0), (0, 13), (0, -13)

iii)



4. a)  $x^2 + y^2 = 121$   
 b)  $x^2 + y^2 = 81$   
 5. a) 17 units  
 b)  $x^2 + y^2 = 289$

- c)  $x^2 + y^2 = 16$   
 d)  $x^2 + y^2 = 36$   
 c)




6. a) yes,  $(-4)^2 + 7^2 = 65$   
 b) no,  $5^2 + (-6)^2 \neq 65$

7. a) i) 5 units  
 ii) 5 units  
 b) i)  $x^2 + y^2 = 25$   
 ii)  $x^2 + y^2 = 25$

c) Answers may vary, e.g.,

- i) (3, 4), (-3, -4)  
 ii) (0, 5), (-5, 0)

d) i)




- c) yes,  $8^2 + (-1)^2 = 65$   
 d) no,  $(-3)^2 + (-6)^2 \neq 65$


- iii) 3 units  
 iv) 17 units  
 iii)  $x^2 + y^2 = 9$   
 iv)  $x^2 + y^2 = 289$

- iii) (3, 0), (-3, 0)  
 iv) (8, 15), (-8, -15)


ii)



iii)



iv)



8. a)  $x^2 + y^2 = 20\,736$     c)  $x^2 + y^2 = 1190.25$   
 b)  $x^2 + y^2 = 361\,000\,000$     d)  $x^2 + y^2 = 1.44$   
 9.  $x^2 + y^2 = 9$ ,  $x^2 + y^2 = 20.25$ ,  $x^2 + y^2 = 56.25$ ,  $x^2 + y^2 = 81$ ,  
 $x^2 + y^2 = 144$   
 10. about 1257 km  
 11. a)  $x^2 + y^2 = 169$     b) (-5, 12)  
 12.  $x^2 + y^2 = 289$   
 13. about 37s  
 14.  $a = 10.0$  or  $-10.0$ ,  $b = 6.6$  or  $-6.6$   
 15. Answers may vary, e.g., I would calculate the distance from (0, 0) to (12 504, 16 050) and compare it with the square root of 45 000 000, which is the radius of the first satellite's orbit.  
 16. about 11.3 units by about 11.3 units  
 17. Answers may vary, e.g.,

Reason 1: The distance from the origin (0, 0) to the point (x, y) is  $\sqrt{(x-0)^2 + (y-0)^2}$ , which is equal to  $\sqrt{x^2 + y^2}$ . If, however, (x, y) is a point on a circle with centre (0, 0) and radius r, then the distance from (0, 0) to (x, y) is r. So  $\sqrt{x^2 + y^2} = r$  and, if you square both sides,  $x^2 + y^2 = r^2$ .

Reason 2: The equation  $x^2 + y^2 = r^2$  follows directly from the Pythagorean theorem applied to the right triangle with vertices at (0, 0), (x, 0), and (x, y).

Reason 3: Using the formula  $x^2 + y^2 = r^2$ , I can see that the x- and y-intercepts are (r, 0), (-r, 0), (0, r), (0, -r). This makes sense for a circle, since all intercepts should be the same distance from the centre.


18. a) a circle centred at the origin, with a radius of  $\frac{4}{3}$   
 b) a circle centred at (2, -4), with a radius of 3  
 19. The most likely part of the load to hit the edge of the tunnel is the corner. The distance from the middle of the road to the corner of the load is  $\sqrt{(4)^2 + (3.5)^2} \approx 5.32$ , or about 5.32 m. Since this is larger than the radius of the tunnel, the load will not fit through the tunnel.

### Mid-Chapter Review, page 95


1. a) (-4, 4)    b) (1.5, 4)    c) (0, 4)    d) (3, 2)  
 2. (6, -3)  
 3.  $y = -2x + 3$   
 4. all points on the equation  $y = \frac{7}{2}x - \frac{39}{4}$   
 5. a)  $M_{PQ} = (3, 3)$ ,  $M_{QR} = (-5, 0)$ ,  $M_{RP} = (4, 1)$   
 b)  $y = -\frac{1}{10}x + \frac{7}{5}$   
 c)  $y = -9x + 30$   
 6. a) about 5.4 units    c) 7 units  
 b) about 12.1 units    d) about 3.2 units  
 7. about 67.2 m  
 8. about 6.3 units

9. length  $AB \cong 5.8$  units, length  $BC \cong 8.2$  units, length  $CA \cong 8.6$  units; so the sides are unequal


10. a) i)  $(0, 0)$   
 ii) radius: 13 units;  $x$ -intercepts:  $(13, 0)$ ,  $(-13, 0)$ ;  
 $y$ -intercepts:  $(0, 13)$ ,  $(0, -13)$



- b) i)  $(0, 0)$   
 ii) radius: 1.7 units;  
 $x$ -intercepts:  $(1.7, 0)$ ,  $(-1.7, 0)$ ;  
 $y$ -intercepts:  $(0, 1.7)$ ,  $(0, -1.7)$



- c) i)  $(0, 0)$   
 ii) radius: about 9.9 units;  
 $x$ -intercepts: about  $(9.9, 0)$ , about  $(-9.9, 0)$ ;  
 $y$ -intercepts: about  $(0, 9.9)$ , about  $(0, -9.9)$



11. a)  $x^2 + y^2 = 25$   
 b)  $x^2 + y^2 = 49$   
 c)  $x^2 + y^2 = 73$   
 d)  $x^2 + y^2 = 97$

12.  $x^2 + y^2 = 900$


13. a) on:  $6^2 + (-3)^2 = 45$   
 b) outside:  $(-1)^2 + 7^2 > 45$   
 c) inside:  $(-3)^2 + 5^2 < 45$   
 d) outside:  $(-7)^2 + (-2)^2 > 45$
14. a)  $6^2 + (-7)^2 = 85$ ,  $2^2 + 9^2 = 85$

b)  $y = \frac{1}{4}x$


- c) because  $(0, 0)$  is the centre of the circle and because  $0 = \frac{1}{4}(0)$ ,  
 $(0, 0)$  also lies on the perpendicular bisector

## Lesson 2.4, page 101


- The slopes are equal:  $m_{PQ} = m_{RS} = \frac{1}{4}$
- The slopes are negative reciprocals:  $m_{TU} = -\frac{1}{m_{VW}} = -\frac{1}{2}$
- $ABCD$  is a parallelogram. The lengths of the sides are needed to determine if it is also a rhombus.
- a)  $DE = FD = \sqrt{65} \cong 8.06$  units;  $EF = \sqrt{130} \cong 11.40$  units  
 b)  $\sqrt{32.5} \cong 5.72$  units  
 c)  $m_{MD} = -\frac{1}{m_{EF}} = \frac{7}{9}$
- $PQRS$  is a parallelogram and a rhombus. The slopes of the sides or the interior angles are needed to determine if it is also a square.
- a) isosceles;  $AB = \sqrt{17} \cong 4.12$  units,  $BC = \sqrt{17} \cong 4.12$  units,  $CA = \sqrt{34} \cong 5.83$  units; two equal sides




- b) scalene;  $GH = \sqrt{26} \cong 5.10$  units,  $HI = \sqrt{20} \cong 4.47$  units,  
 $GI = \sqrt{18} \cong 4.24$  units; no equal sides



- c) scalene;  $DE = \sqrt{17} \cong 4.12$  units,  $EF = \sqrt{68} \cong 8.25$  units,  
 $DF = 5$  units; no equal sides




- d) isosceles;  $JK = \sqrt{58} \cong 7.62$  units,  $KL = 6$  units,  
 $LJ = \sqrt{58} \cong 7.62$  units; two equal sides




7.  $m_{PQ} = -\frac{1}{m_{PR}} = -3$ , so  $PQ$  is perpendicular to  $PR$ ; that is,  $PQ$  meets  $PR$  at a right angle.
8.  $m_{MN} = -\frac{1}{m_{NL}} = -\frac{4}{5}$ ;  $MN = NL = \sqrt{41} \doteq 6.40$  units
9. a) Use the distance formula to determine the lengths of the three sides. If the lengths make the equation  $a^2 + b^2 = c^2$  true, then the triangle is a right triangle.  
 b) i)  $\triangle STU$  is a right triangle because  $ST^2 + TU^2 = 17 + 68 = 85 = US^2$   
 ii)  $\triangle XYZ$  is not a right triangle because  $XY^2 = 20$ ,  $YZ^2 = 80$ , and  $XZ^2 = 52$   
 iii)  $\triangle ABC$  is a right triangle because  $AB^2 + AC^2 = 13 + 13 = 26 = BC^2$
10. a)  $WX = \sqrt{29} \doteq 5.39$  units,  $XY = \sqrt{41} \doteq 6.40$  units,  $YZ = \sqrt{29} \doteq 5.39$  units,  $ZW = \sqrt{41} \doteq 6.40$  units;  $m_{WX} = 0.4$ ,  $m_{XY} = -1.25$ ,  $m_{YZ} = 0.4$ ,  $m_{ZW} = -1.25$   
 b) parallelogram; because opposite sides are equal length and parallel (since they have the same slope)  
 c)  $\sqrt{90} - \sqrt{50} \doteq 2.42$  units
11.  $m_{RS} = m_{TU} = \frac{2}{10}$ ,  $m_{ST} = m_{UR} = \frac{4}{3}$  or  $RS = TU = \sqrt{104} \doteq 10.20$ ,  $ST = UR = 5$ , so opposite sides are equal length and parallel (because they have the same slope).
12.  $AB = BC = CD = DA = 5$ , so all sides are equal.
13. a)  $EF = FG = GH = HE = \sqrt{20} \doteq 4.47$ , so all sides are equal;  $m_{EF} = m_{GH} = -\frac{1}{2}$ ,  $m_{FG} = m_{EH} = 2$ , so adjacent sides meet at right angles.  
 b)  $m_{EG} = -3$ ,  $m_{HF} = \frac{1}{3}$ ; the slopes of  $EG$  and  $HF$  are negative reciprocals, so  $EG$  and  $HF$  are perpendicular to each other.
14.  $m_{PQ} = \frac{7}{9}$ ,  $m_{QR} = \frac{3}{2}$ ;  $m_{PQ} \neq -\frac{1}{m_{QR}}$ ; the slopes are not negative reciprocals, so the sides are not perpendicular; that is, they do not meet at right angles.
15. Answers may vary, e.g., I would use the distance formula to determine the lengths of all the sides. If they are equal, then the quadrilateral is a rhombus; e.g.,  $A(3, 0)$ ,  $B(0, 2)$ ,  $C(-3, 0)$ ,  $D(0, -2)$ . Or, I would use the slope formula to determine the slopes of all the sides. If the slopes of adjacent sides are negative reciprocals of each other, then the quadrilateral is a rectangle, e.g.,  $E(0, 0)$ ,  $F(2, 1)$ ,  $G(0, 5)$ ,  $H(-2, 4)$ . Or, if the quadrilateral is both a rhombus and a rectangle, then it is a square; e.g.,  $J(3, 0)$ ,  $K(0, 4)$ ,  $L(-4, 1)$ ,  $M(-1, -3)$ .
16. a) rhombus;  $JK = KL = LM = JM = \sqrt{17} \doteq 4.12$  units;  $m_{JK} = m_{LM} = \frac{1}{4}$ ,  $m_{KL} = m_{JM} = 4$ ; all sides are equal length, but there are no right angles.  
 b) parallelogram;  $EF = GH = 5$ ,  $FG = EH = \sqrt{153} \doteq 12.37$  units;  $m_{FG}$  and  $m_{EH}$  are undefined (vertical),  $m_{FG} = m_{EH} = \frac{1}{4}$ , opposite sides are equal length and parallel, but there are no right angles.  
 c) parallelogram;  $DE = FG = \sqrt{50} \doteq 7.07$  units,  $EF = GH = \sqrt{29} \doteq 5.39$  units;  $m_{DE} = m_{FG} = \frac{1}{7}$ ,  $m_{EF} = m_{DG} = \frac{5}{2}$ ; opposite sides are equal length and parallel, but there are no right angles.

- d) rectangle;  $PQ = RS = \sqrt{68} \doteq 8.25$  units,  $QR = PS = \sqrt{17} \doteq 4.12$  units;  $m_{PQ} = m_{RS} = \frac{1}{4}$ ,  $m_{QR} = m_{PS} = -4$ ; opposite sides are equal length and parallel, and angles between sides are  $90^\circ$ .
17. square; all side lengths are  $\sqrt{106} \doteq 10.30$ , or about 10.30 units, so the side lengths are equal; slopes are  $\frac{5}{9}$ ,  $-\frac{9}{5}$ ,  $\frac{5}{9}$ ,  $-\frac{9}{5}$ , so the slopes of the sides are negative reciprocals.
18. a)  $S(8, 2)$   
 b) Answers may vary, e.g., I determined the difference of the  $x$ - and  $y$ -coordinates between  $Q$  and  $P$  and then applied this difference to  $R$ .  
 c) yes,  $PR = QS = \sqrt{145} \doteq 12.04$ , or about 12.04 units
19. Answers may vary, e.g.,




20. Answers may vary, e.g.,



Since  $\frac{AC}{FC} = \frac{BC}{EC}$  and  $\angle ACB = \angle FCE$ ,  $\triangle ABC$  is similar to  $\triangle FEC$ . Each side in the larger triangle is twice the length of the corresponding side in the smaller triangle.

b)




$\triangle AHE$  is similar to  $\triangle ADB$ , so  $HE \parallel DB$ .  
 $\triangle CGF$  is similar to  $\triangle CDB$ , so  $GF \parallel DB$ .  
 $\triangle DGH$  is similar to  $\triangle DCA$ , so  $GH \parallel CA$ .  
 $\triangle BFE$  is similar to  $\triangle BCA$ , so  $FE \parallel CA$ .  
 Since  $HE \parallel DB \parallel GF$  and  $GH \parallel CA \parallel FE$ ,  $EFGH$  is a parallelogram.

### Lesson 2.5, page 109

- $AC = BD = \sqrt{65} \approx 8.06$ , so  $AC$  and  $BD$  are each about 8.06 units.
- $m_{JL} = -\frac{1}{2}$ ,  $m_{KM} = 2$ ; the slopes of the diagonals are negative reciprocals, so the diagonals are perpendicular.
- Let  $S$  represent the midpoint of  $PR$ . Since  $S = (1.5, 1.5)$ ,  $QS$  bisects  $PR$ . Since  $m_{QS} = -\frac{1}{m_{PR}} = -7$ , they are perpendicular.
- $M_{JL} = M_{KM} = (-1, -3)$
- $AC = BD = \sqrt{520} \approx 22.80$  units
- Answers may vary, e.g., conjecture: Quadrilateral  $ABCD$  is a rectangle. I calculated the length and slope of each blue line segment and found that opposite sides are equal, and adjacent sides are perpendicular. My conjecture was correct.
- Answers may vary, e.g., conjecture: Quadrilateral  $JKLM$  is a kite. I calculated the length of each blue line segment and found that adjacent sides  $JK$  and  $KL$  are equal. Also adjacent sides  $JM$  and  $ML$  are equal. My conjecture was correct.
- Let  $G$  represent the midpoint of  $EF$ ;  $G = (0, 3)$ . Since  $m_{EF} = -\frac{1}{m_{DG}} = 5$ ,  $DG$  is the median and the altitude.
- Answers may vary, e.g.,  $M_{PQ} = (-1, 1)$ ,  $M_{QR} = (3, 3.5)$ ,  $M_{RS} = (7, 1)$ ,  $M_{SP} = (3, -1.5)$ ;  $M_{PQ}M_{QR} = M_{QR}M_{RS} = M_{RS}M_{SP} = M_{SP}M_{PQ} = \sqrt{22.25} \approx 4.72$ , or about 4.72 units; therefore,  $M_{PQ}M_{QR}M_{RS}M_{SP}$  is a rhombus.
- Answers may vary, e.g.,  $M_{RS} = (-3, 2.5)$ ,  $M_{ST} = (-1.5, 1)$ ,  $M_{TU} = (-4, -1.5)$ ,  $M_{UR} = (-5.5, 0)$ ; diagonals  $M_{RS}M_{TU} = M_{ST}M_{UR} = \sqrt{17} \approx 4.12$ , or about 4.12 units; therefore, the midpoints of the rhombus form a rectangle.
- Answers may vary, e.g.,  $m_{RT} = -\frac{1}{m_{SU}} = -1$ ; the slopes of the diagonals are negative reciprocals, so the diagonals are perpendicular.  $M_{RT} = M_{SU} = (-3.5, 0.5)$ , so the diagonals bisect each other.
- Answers may vary, e.g.,  $M_{AB} = (-4, -10)$ ,  $M_{BC} = (-8, -2)$ ,  $M_{CD} = (0, 2)$ ,  $M_{DA} = (4, -6)$ ;  $M_{AB}M_{BC} = M_{BC}M_{CD} = M_{CD}M_{DA} = M_{DA}M_{AB} = \sqrt{80} \approx 8.94$  units, so the midsegments form a rhombus.  $M_{AB}M_{CD} = M_{BC}M_{DA} = \sqrt{160} \approx 12.65$ , or about 12.65 units, so the rhombus is a rectangle and, therefore, a square.


- a)  $(-4)^2 + 3^2 = 25$ ,  $3^2 + (-4)^2 = 25$   
 b)  $m_{AB} = -1$ ; therefore, the equation of the perpendicular bisector of  $AB$  is  $y = x$ ; for  $y = x$ , when  $x = 0$  and  $y = 0$ , left side equals 0 and right side equals 0 so the centre  $(0, 0)$  of the circle lies on the perpendicular bisector.
- a)  $M_{BC} = (-3, -0.5)$ ,  $M_{AD} = (1.5, 1)$ ; slope of  $M_{BC}M_{AD} = m_{AB} = m_{DC} = \frac{1}{3}$ ; the slopes are the same, so the line segments are parallel.  
 b)  $M_{BC}M_{AD} = \sqrt{22.5} \approx 4.74$ , or about 4.74 units;  
 $\frac{BC + AD}{2} = \frac{\sqrt{10} + \sqrt{40}}{2} = 1.5\sqrt{10} = \sqrt{22.5} \approx 4.74$ , or about 4.74 units
- Answers may vary, e.g., area of  $\triangle ABC = \frac{1}{2}(7)(4) = 14$ , or 14 square units, area of  $\triangle M_{AB}M_{BC}M_{AC} = \frac{1}{2}(3.5)(2) = 3.5 = \frac{1}{4}(14)$ , or one-quarter of 14 square units
- Answers may vary, e.g.,



- Answers may vary, e.g., let the vertices of the square be  $A(0, 0)$ ,  $B(2a, 0)$ ,  $C(2a, 2a)$ , and  $D(0, 2a)$ . The midpoints of  $M_{AB}M_{CD}$ ,  $M_{BC}M_{AD}$ ,  $AC$ , and  $BD$  are all  $(a, a)$ .

### Lesson 2.6, page 113

- $(1, 4)$
- Answers may vary, e.g.,



Let  $AD$  represent the median, and let  $AE$  represent the altitude. Both  $\triangle ACD$  and  $\triangle ABD$  have the same base (since  $CD = DB$ ) and the same height,  $AE$ . Therefore, they have the same area.

- a)  $AE \times EB = CE \times ED = 60$   
 b) 1.25 m
- a)



b) Answers may vary, e.g.,

Similar Figures	Diagram	$A_1$ : Area on 6 cm Side ( $\text{cm}^2$ )	$A_2$ : Area on 8 cm Side ( $\text{cm}^2$ )	$A_1 + A_2$ ( $\text{cm}^2$ )	Area on Hypotenuse ( $\text{cm}^2$ )
square		36	64	100	100
semicircle		14.14	25.13	39.27	39.27
rectangle		18	32	50	50
equilateral triangle		15.59	27.71	43.30	43.30
right triangle		9	16	25	25
parallelogram		18	32	50	50

c) no effect

d) The sum of the two smaller areas always equals the larger area.

### Lesson 2.7, page 120

- $\frac{1}{2}$
  - $-2$
  - $y = -2x + 2$
- $y = \frac{1}{2}x - \frac{3}{2}$
- $\left(\frac{7}{5}, \frac{4}{5}\right)$
- $BC = \sqrt{45} \doteq 6.71$ , or about 6.71 units;  
 $AD = \sqrt{28.8} \doteq 5.37$ , or about 5.37 units
- 18 square units
- $y = -\frac{5}{3}x - 3$
  - $y = -\frac{5}{3}x - 3$
  - $y = -\frac{5}{3}x - 3$
- isosceles; the median and the altitude are the same.

7. a)  $x^2 + y^2 = 85$

b)  $7^2 + 6^2 = 85$

c)  $m_{PR} = -\frac{1}{m_{RQ}} = \frac{1}{4}$ , so the slopes are negative reciprocals;  
therefore,  $\angle PRQ$  is a right angle.

8. about 27 square units

9.  $\left(2, \frac{16}{3}\right)$

10.  $\left(-\frac{16}{3}, 0\right)$

11. (2, 3)

12. (6, 4)

13. Answers may vary, e.g.,

i) Calculate  $DC$ ,  $EC$ , and  $FC$ ; check that they are equal.

ii) Construct the perpendicular bisectors for  $\triangle DEF$ ; check that their intersection is at  $C$ .

14.  $\left(\frac{22}{3}, 8\right)$


15. 3:1

16. a) House  $A$ : about 37.2 m; house  $B$ : about 26.8 m; house  $C$ : about 34.8 m; assuming that the connection charge is proportional to distance,  $A$  has the highest charge.

b) about (18.1, 14.2)



17. right triangle; the lines forming two of the sides are perpendicular, having slopes  $-1$  and  $1$ .



Use the chord intersection rule to find  $E$  such that  $AD \times DC = BD \times DE$ ;  $BD + DE = \text{diameter}$ .

19. Answers may vary, e.g., to determine the median, I would calculate the midpoint of the opposite side and determine the slope of the segment between the vertex and the midpoint, which gives  $m$ . Then I would substitute one of these points into the equation of a line to determine  $b$ . To determine the altitude, I would calculate the slope of the opposite side and determine its negative reciprocal, which gives  $m$ . Then I would substitute the vertex into the equation of a line to determine  $b$ .
20. Let  $C$  represent the centroid;  
 $M_{PQ} = \left(\frac{3}{2}, -1\right)$ ,  $M_{QR} = \left(\frac{5}{2}, -1\right)$ ,  $M_{RP} = (0, 2)$ ,  $C = \left(\frac{4}{3}, 0\right)$   
 $\frac{PC}{CM_{QR}} = \frac{QC}{CM_{RP}} = \frac{RC}{CM_{PQ}} = 2$
21. a)  $(3a)^2 + a^2 = 10a^2$ ,  $a^2 + (-3a)^2 = 10a^2$   
 b) Let  $C$  represent the centre;  $C = (0, 0)$ ,  $M_{RQ} = (2a, -a)$ ; slope  $RQ = 2$ , slope  $CM_{RQ} = -\frac{1}{2}$ ; the slopes are negative reciprocals. Therefore, the line segments are perpendicular.

## Chapter Review, page 124

- $(7.5, 28)$
- a)  $y = x + 2$   
 b) yes,  $m_{\text{median}} = -\frac{1}{m_{AC}} = 1$
- $y = \frac{7}{4}x + \frac{21}{8}$
- $Q$
- about 89.5 units
- about 21.8 units
- 10 units
- a)  $x^2 + y^2 = 289$   
 b)  $x$ -intercepts:  $(17, 0)$ ,  $(-17, 0)$ ;  $y$ -intercepts:  $(0, 17)$ ,  $(0, -17)$ ; points: e.g.,  $(8, 15)$ ,  $(8, -15)$ ,  $(-8, -15)$
- $x^2 + y^2 = 841$
- $x^2 + y^2 = 16$
- $(-2)^2 + k^2 = 20$   
 $k^2 = 16$   
 $k = 4$  or  $-4$
- $AB = BC = \sqrt{17} \approx 4.12$ , or about 4.12 units; two sides are equal in length, so the triangle is isosceles.
- $AB = \sqrt{13} \approx 3.61$ ,  $BC = \sqrt{26} \approx 5.10$ ,  $CA = \sqrt{13} \approx 3.61$ , or about 3.61 units; two sides are equal in length, so the triangle is isosceles.
- $JK = KL = LM = MJ = 5$  units; the sides are equal in length, so the quadrilateral is a rhombus.

15.  $RS = TU = \sqrt{20} \approx 4.47$ ,  $ST = RU = \sqrt{17} \approx 4.12$ , or about 4.12 units

$$m_{RS} = 2, m_{ST} = -\frac{1}{4}, m_{TU} = 2, m_{UR} = -\frac{1}{4}$$

The opposite sides are equal and parallel, but the adjacent sides do not meet at  $90^\circ$ . Therefore, the quadrilateral is a parallelogram.

16.  $M_{AB} = (-4, -4)$ ,  $M_{BC} = (1, -5)$ ,  $M_{CD} = (5, 1)$ ,  $M_{DA} = (0, 2)$ ;  
 $M_{AB}M_{BC} = M_{CD}M_{DA} = \sqrt{26} \approx 5.10$ , or about 5.10 units;  
 $M_{BC}M_{CD} = M_{DA}M_{AB} = \sqrt{52} \approx 7.21$ , or about 7.21 units;  
 The slopes of the midsegments are  $\frac{3}{2}$  and  $-\frac{1}{5}$ . The opposite sides are equal and parallel, but the adjacent sides do not meet at  $90^\circ$ . Therefore, the quadrilateral is a parallelogram.

17.  $(10 - 5)^2 + (10 + 2)^2 = (-7 - 5)^2 + (3 + 2)^2 = (0 - 5)^2 + (-14 + 2)^2$

18. a)  $m_{PQ} = -\frac{1}{m_{QR}} = \frac{5}{2}$ ; the slopes of  $PQ$  and  $QR$  are negative reciprocals, so  $PQ$  and  $QR$  form a right angle.

b)  $M_{RP} = \left(2, \frac{5}{2}\right)$ ;  $PM_{RP} = QM_{RP} = RM_{RP} = \sqrt{36.25} \approx 6.02$ , or about 6.02 units

19. a)  $6^2 + 7^2 = (-9)^2 + 2^2$ , so both points are the same distance from  $(0, 0)$ .

- b) Let  $A$  represent point  $(6, 7)$ . Let  $B$  represent point  $(-9, 2)$ . Let  $C$  represent the centre of the circle  $(0, 0)$ . Let  $D$  represent the

intersection of the line and the chord.  $M_{AB} = \left(-\frac{3}{2}, \frac{9}{2}\right)$ ;

$$m_{AB} = \frac{1}{3}; m_{CD} = -3; \text{ therefore, the equation of the line through}$$

$C$  is  $y = -3x$ . Since  $M_{AB}$  is on this line,  $D = M_{AB}$ .

20. a)  $M_{JL} = M_{KM} = (5.5, -4.5)$   
 b) Answers may vary, e.g., conjecture: Quadrilateral  $JKLM$  is a square.  
 c) Answers may vary, e.g.,  $JK = KL = LM = MJ = \sqrt{13} \approx 3.61$ , or about 3.61 units; I calculated to determine that opposite sides are equal and that adjacent sides are perpendicular. My conjecture was correct.

21.  $(0.5, 2)$

22.  $(3, 2)$

23. about  $(19.9, 89.3)$

24. Answers may vary, e.g., it will be a parallelogram because the slopes of the lines form two pairs, which are not negative reciprocals;

$$m_1 = -3, m_2 = \frac{4}{5}, m_3 = -3, m_4 = \frac{4}{5}$$

25. a)  $\left(\frac{453}{17}, \frac{194}{17}\right)$

- b) about 6.79 m

## Chapter Self-Test, page 126

- a) about 40.3 m  
 b)  $\left(4, \frac{11}{2}\right)$
- a)  $x^2 + y^2 = 1296$   
 b) 5 s