




## LESSON 77 - Geometric Representations of Complex Numbers

Argand Diagram  
Modulus and Argument  
Polar form

### Argand Diagram

- Complex numbers can be shown Geometrically on an Argand diagram
- The real part of the number is represented on the x-axis and the imaginary part on the y.

- 3
- 4i
- $3 + 2i$
- $2 - 2i$



### Modulus of a complex number


- A complex number can be represented by the position vector.

$$\begin{pmatrix} x \\ y \end{pmatrix}$$

- The Modulus of a complex number is the distance from the origin to the point.

Can you generalise this?

$$|z| = \sqrt{x^2 + y^2}$$



How many complex numbers in the form  $a + bi$  can you find with integer values of  $a$  and  $b$  that share the same modulus as the number above.

Could you mark all of the points?

What familiar shape would you draw?(more of LOCI later!)



### Geogebra Illustrations of Algebra of Complex Numbers

<http://tube.geogebra.org/material/simple/id/23937>

<http://tube.geogebra.org/material/simple/id/23938>

<http://tube.geogebra.org/material/simple/id/42722>

● ● ● | Modulus questions

Find

a)  $|3 + 4i|$

b)  $|5 - 12i|$

c)  $|6 - 8i|$

d)  $|-24 - 10i|$

Find the distance between the first two complex numbers above. It may help to sketch a diagram

● ● ● | Example

**EXAMPLE 11.16**

If  $z = 1 + 2i$  and  $w = x - i$ , find

(a)  $|z + 4|$

(b)  $|z + w|$

Find the minimum value of  $|z + w|$ .

● ● ● | Example - Solution

**Solution**

(a) First, we need to determine the complex number  $z + 4$ :

$$z + 4 = (1 + 2i) + 4 = 5 + 2i.$$

$$\text{Then we have, } |5 + 2i| = \sqrt{25 + 4} = \sqrt{29}.$$

(b) First, we need to determine the complex number  $z + w$ :

$$z + w = (1 + 2i) + (x - i) = (x + 1) + i \quad \therefore |(x + 1) + i| = \sqrt{(x + 1)^2 + 1}.$$

$$= \sqrt{x^2 + 2x + 2}$$

395

**MATHEMATICS – Higher Level (Core)**

Now,  $\sqrt{x^2 + 2x + 2} = \sqrt{(x^2 + 2x + 1) + 1} = \sqrt{(x + 1)^2 + 1}$ . The minimum value will occur when  $(x + 1)^2 + 1$  is a minimum. But, the minimum value of  $(x + 1)^2 + 1$  is 1, therefore the minimum of  $\sqrt{(x + 1)^2 + 1}$  is  $\sqrt{1} = 1$ .

● ● ● | The argument of a complex number


The argument of a complex number is the angle the line makes with the positive x-axis.

Can you generalise this?

$$\arg z = \theta$$

$$\theta = \tan^{-1}(y / x)$$

$$-\pi \leq \theta \leq \pi$$



It is really important that you sketch a diagram before working out the argument!!

### ● ● ● | The argument of a complex number

- Calculate the modulus and argument of the following complex numbers. (Hint, it helps to draw a diagram)

1)  $3 + 4i$

2)  $5 - 5i$

3)  $-2\sqrt{3} + 2i$

### ● ● ● | The Polar form of a complex number

- So far we have plotted the position of a complex number on the Argand diagram by going horizontally on the real axis and vertically on the imaginary.

- This is just like plotting co-ordinates on an x,y axis

- However it is also possible to locate the position of a complex number by the distance travelled from the origin (pole), and the angle turned through from the positive x-axis.



- These are called "Polar coordinates"

### ● ● ● | The Polar form of a complex number

$(x, y)$   $r$  is the MODULUS  $(r, \theta)$  The ARGUMENT  
 $\cos\theta = x/r, \sin\theta = y/r$   
 $x = r \cos\theta, y = r \sin\theta$

REAL Part

IMAGINARY part



### ● ● ● | Converting from Cartesian to Polar


$$(x, y) = (r, \theta) = \left( \sqrt{x^2 + y^2}, \tan^{-1} \frac{y}{x} \right)$$

Convert the following from Cartesian to Polar

i)  $(1, 1) =$

ii)  $(-\sqrt{3}, 1) =$

iii)  $(-4, -4\sqrt{3}) =$



### Converting from Cartesian to Polar


$$(x, y) = (r, \theta) = \left( \sqrt{x^2 + y^2}, \tan^{-1} \frac{y}{x} \right)$$

Convert the following from Cartesian to Polar

i)  $(1, 1) = (\sqrt{2}, \pi/4)$

ii)  $(-\sqrt{3}, 1) = (2, 5\pi/6)$

iii)  $(-4, -4\sqrt{3}) = (8, -2\pi/3)$



### Converting from Polar to Cartesian


$$(r, \theta) = (x, y) = (r \cos \theta, r \sin \theta)$$

Convert the following from Polar to Cartesian

i)  $(4, \pi/3) =$

ii)  $(3\sqrt{2}, -\pi/4) =$

iii)  $(6\sqrt{2}, 3\pi/4) =$



### Converting from Polar to Cartesian


$$(r, \theta) = (x, y) = (r \cos \theta, r \sin \theta)$$

Convert the following from Polar to Cartesian

i)  $(4, \pi/3) = (2, 2\sqrt{3})$


ii)  $(3\sqrt{2}, -\pi/4) = (3, -3)$

iii)  $(6\sqrt{2}, 3\pi/4) = (-6, 6)$




If  $r \geq 0$  and  $0 \leq \theta < 2\pi$ , the complex number  $z = x + yi$  may be written in polar form as

$$z = x + yi = (r \cos \theta) + (r \sin \theta)i = r(\cos \theta + i \sin \theta) \quad (4)$$



- Plot the point corresponding to  $z = -3 + 4i$  in the complex plane, and write an expression for  $z$  in polar form.


- Plot the point corresponding to  $z = -3 + 4i$  in the complex plane, and write an expression for  $z$  in polar form.







- $z = -3 + 4i$  is in Quadrant II


$$x = -3 \text{ and } y = 4$$

$$r = \sqrt{(-3)^2 + 4^2} = \sqrt{25} = 5$$






Find the reference angle ( $\alpha$ ) by solving



$$\tan \alpha = \left| \frac{y}{x} \right|$$

$$\tan \alpha = \left| \frac{-1}{\sqrt{3}} \right|$$

$$\alpha = \tan^{-1} \left| \frac{-1}{\sqrt{3}} \right|$$

$$\alpha = 30^\circ$$


$$\theta = 360^\circ - 30^\circ = 330^\circ$$

$$z = r(\cos \theta + i \sin \theta)$$

$$z = 2(\cos 330^\circ + i \sin 330^\circ)$$

Write in standard (rectangular) form.

$$2 \left( \cos \frac{5\pi}{6} + i \sin \frac{5\pi}{6} \right)$$

Write in standard (rectangular) form.

$$2 \left( \cos \frac{5\pi}{6} + i \sin \frac{5\pi}{6} \right)$$

$$\cos \frac{5\pi}{6} = -\frac{\sqrt{3}}{2} \quad \sin \frac{5\pi}{6} = \frac{1}{2}$$

$$2 \left( -\frac{\sqrt{3}}{2} + \frac{1}{2}i \right) = -\sqrt{3} + i$$

Example:

Convert  $z = 3 \operatorname{cis} 55^\circ$  to rectangular form.

Convert  $z = -2 - 3i$  to polar form.

Example:

What is the absolute value of the following complex numbers:

$$z = 3 + 2i$$

$$z = 4 \operatorname{cis} \frac{2\pi}{3}$$

Multiply:  $(3 \operatorname{cis} 165^\circ)(4 \operatorname{cis} 45^\circ)$

Do you want to go thru that every time?

No!

$$(r \operatorname{cis} \alpha) \cdot (t \operatorname{cis} \beta) = r \cdot t \cdot \operatorname{cis} (\alpha + \beta)$$



**Multiply:**  $(4\text{cis}25^\circ)(6\text{cis}35^\circ)$

**Divide:**  $\frac{3\text{cis}165^\circ}{4\text{cis}45^\circ}$

### Properties of modulus

$$|z_1 \cdot z_2| = |z_1| \cdot |z_2| \quad |z^n| = |z|^n$$

$$|\bar{z}| = |z|$$

$$|z \cdot \bar{z}| = |z|^2 \quad \text{Proof: } z = a + ib, \bar{z} = (a - ib)(a - ib) = a^2 + b^2 = |z|^2$$

$$\left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|}$$

$$|z_1 + z_2| \leq |z_1| + |z_2| \quad |z_1 + z_2| \geq ||z_1| - |z_2|| \quad (\text{Triangle inequality})$$

$$|z_1 - z_2| \geq ||z_1| - |z_2|| \quad |z_1 - z_2| \leq |z_1| + |z_2|$$

$$|z_1 - z_2|^2 + |z_1 + z_2|^2 = 2(|z_1|^2 + |z_2|^2)$$

### Properties of Argument

$$\text{Arg}(z_1 z_2) = \text{Arg}(z_1) + \text{Arg}(z_2)$$

$$\text{Arg}\left(\frac{z_1}{z_2}\right) = \text{Arg}(z_1) - \text{Arg}(z_2)$$

$$\text{Arg}(z_1 z_2 \dots z_n) = \text{Arg}(z_1) + \text{Arg}(z_2) + \dots + \text{Arg}(z_n)$$

$$\text{Arg}(\bar{z}) = -\text{Arg}(z), \quad \text{Arg}\left(\frac{1}{z}\right) = -\text{Arg}(z)$$

**Arg(purely real) = 0 or  $\pi$  or  $2n\pi$  and vice versa**

**Arg(purely imaginary) =  $\frac{\pi}{2}$  or  $-\frac{\pi}{2}$  or  $(2n+1)\frac{\pi}{2}$  and vice versa**