

SOLVED PROBLEMS ON TAYLOR AND MACLAURIN SERIES

TAYLOR AND MACLAURIN SERIES

Taylor Series of a function f at $x = a$ is

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (x - a)^k$$

It is a Power Series centered at a .

Maclaurin Series of a function f is a Taylor Series at $x = 0$.

BASIC MACLAURIN SERIES

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots = \sum_{k=0}^{\infty} \frac{x^k}{k!}$$

$$\sin(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots = \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{(2k+1)!}$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots = \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k}}{(2k)!}$$

$$(1+x)^p = 1 + px + \frac{p(p-1)}{2!} x^2 + \dots$$

USE TAYLOR SERIES

- 1 To estimate values of functions on an interval.
- 2 To compute limits of functions.
- 3 To approximate integrals.
- 4 To study properties of the function in question.

FINDING TAYLOR SERIES

To find Taylor series of functions, we may:

- 1 Use substitution.
- 2 Differentiate known series term by term.
- 3 Integrate known series term by term.
- 4 Add, divide, and multiply known series.

OVERVIEW OF PROBLEMS

Find the Maclaurin Series of the following functions.

- | | | | | | |
|---|-------------|---|---------------------|---|--------------------|
| 1 | $\sin(x^2)$ | 2 | $\frac{\sin(x)}{x}$ | 3 | $\arctan(x)$ |
| 4 | $\cos^2(x)$ | 5 | $x^2 e^x$ | 6 | $\sqrt{1 - x^3}$ |
| 7 | $\sinh(x)$ | 8 | $\frac{e^x}{1 - x}$ | 9 | $x^2 \arctan(x^3)$ |

OVERVIEW OF PROBLEMS

Find the Taylor Series of the following functions at the given value of a .

10 $x - x^3$ at $a = -2$

11 $\frac{1}{x}$ at $a = 2$

12 e^{-2x} at $a = 1/2$

13 $\sin(x)$ at $a = \pi/4$

14 10^x at $a = 1$

15 $\ln(1 + x)$ at $a = -2$

Find the Maclaurin Series of the following functions.

MACLAURIN SERIES

Problem 1

$$f(x) = \sin(x^2)$$

Solution

Substitute x by x^2 in the Maclaurin Series of sine.

$$\text{Hence } \sin(x^2) = \sum_{k=0}^{\infty} (-1)^k \frac{(x^2)^{2k+1}}{(2k+1)!} = \sum_{k=0}^{\infty} (-1)^k \frac{x^{4k+2}}{(2k+1)!}$$

MACLAURIN SERIES

Problem 2

$$f(x) = \frac{\sin(x)}{x}$$

Solution

Divide the Maclaurin Series of sine by x . Hence,

$$\frac{\sin(x)}{x} = \frac{1}{x} \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{(2k+1)!} = \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k}}{(2k+1)!}$$

MACLAURIN SERIES

Problem 3

$$f(x) = \arctan(x)$$

Solution

Observe that $f'(x) = \frac{1}{1+x^2}$. To find the Maclaurin Series of $f'(x)$ substitute $-x^2$ for x in Basic Power Series formula.

MACLAURIN SERIES

Solution(cont'd)

$$\text{Hence } f'(x) = \frac{1}{1+x^2} = \sum_{k=0}^{\infty} (-x^2)^k = \sum_{k=0}^{\infty} (-1)^k x^{2k}.$$

By integrating both sides, we obtain

$$\begin{aligned} f(x) &= \int \left(\sum_{k=0}^{\infty} (-1)^k x^{2k} \right) dx = \sum_{k=0}^{\infty} (-1)^k \int x^{2k} dx \\ &= \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{2k+1} + C. \end{aligned}$$

MACLAURIN SERIES

Solution(cont'd)

0 is in the interval of convergence. Therefore we can insert $x = 0$ to find that the integration constant $c = 0$. Hence the Maclaurin series of $\arctan(x)$ is

$$\arctan(x) = \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{2k+1}.$$

MACLAURIN SERIES

Problem 4

$$f(x) = \cos^2(x)$$

Solution

By the trigonometric identity,

$$\cos^2(x) = (1 + \cos(2x))/2.$$

Therefore we start with the Maclaurin Series of cosine.

MACLAURIN SERIES

Solution(cont'd)

Substitute x by $2x$ in $\cos(x) = \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k}}{(2k)!}$.

Thus $\cos(2x) = \sum_{k=0}^{\infty} (-1)^k \frac{(2x)^{2k}}{(2k)!}$. After adding 1

and dividing by 2, we obtain

MACLAURIN SERIES

Solution(cont'd)

$$\begin{aligned}\cos^2(x) &= \frac{1}{2} \left(1 + \sum_{k=0}^{\infty} (-1)^k \frac{(2x)^{2k}}{(2k)!} \right) \\ &= \frac{1}{2} \left(1 + 1 - \frac{(2x)^2}{2!} + \frac{(2x)^4}{4!} - \dots \right) \\ &= 1 + \sum_{k=1}^{\infty} (-1)^k \frac{2^{2k-1}}{(2k)!} x^{2k}\end{aligned}$$

MACLAURIN SERIES

Problem 5

$$f(x) = x^2 e^x$$

Solution

Multiply the Maclaurin Series of e^x by x^2 .

$$\text{Hence, } x^2 e^x = x^2 \sum_{k=0}^{\infty} \frac{x^k}{k!} = \sum_{k=0}^{\infty} \frac{x^{k+2}}{k!}.$$

MACLAURIN SERIES

Problem 6

$$f(x) = \sqrt{1 - x^3}$$

Solution

By rewriting $f(x) = \left(1 + (-x^3)\right)^{1/2}$. By substituting x by $-x^3$ in the binomial formula with $p = 1/2$ we obtain ,

$$\sqrt{1 - x^3} = 1 - \frac{1}{2} x^3 - \frac{1}{8} x^6 - \dots$$

MACLAURIN SERIES

Problem 7

$$f(x) = \sinh(x)$$

Solution

By rewriting $f(x) = \frac{e^x - e^{-x}}{2}$. Substitute x by $-x$ in

the Maclaurin Series of $e^x = 1 + x + \frac{x^2}{2} + \dots = \sum_{k=0}^{\infty} \frac{x^k}{k!}$,

MACLAURIN SERIES

Solution(cont'd)

$$e^{-x} = \sum_{k=0}^{\infty} \frac{(-x)^k}{k!} = 1 - x + \frac{x^2}{2!} - \dots$$

Thus when we add e^x and e^{-x} , the terms with odd power are canceled and the terms with even power are doubled. After dividing by 2, we obtain

$$\sinh(x) = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \dots = \sum_{k=0}^{\infty} \frac{x^{2k}}{(2k)!}$$

MACLAURIN SERIES

Problem 8

$$f(x) = \frac{e^x}{1-x}$$

Solution

We have $e^x = 1 + x + \frac{x^2}{2!} + \dots$ and $\frac{1}{1-x} = 1 + x + x^2 + \dots$

To find the Maclaurin Series of $f(x)$, we multiply these series and group the terms with the same degree.

MACLAURIN SERIES

Solution(cont'd)

$$\begin{aligned} & \left(1 + x + \frac{x^2}{2!} + \dots \right) \times \left(1 + x + x^2 + \dots \right) \\ &= 1 + 2x + \left(1 + 1 + \frac{1}{2!} \right) x^2 + \text{higher degree terms} \\ &= 1 + 2x + \frac{5}{2} x^2 + \text{higher degree terms} \end{aligned}$$

MACLAURIN SERIES

Problem 9

$$f(x) = x^2 \arctan(x^3)$$

Solution

We have calculated the Maclaurin Series of $\arctan(x)$

$$\arctan(x) = \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{2k+1}.$$

Substituting x by x^3 in the above formula, we obtain

MACLAURIN SERIES

Solution(cont'd)

$$\arctan(x^3) = \sum_{k=0}^{\infty} (-1)^k \frac{(x^3)^{2k+1}}{2k+1} = \sum_{k=0}^{\infty} (-1)^k \frac{x^{6k+3}}{2k+1}.$$

Multiplying by x^2 gives the desired Maclaurin Series

$$x^2 \arctan(x^3) = x^2 \sum_{k=0}^{\infty} (-1)^k \frac{x^{6k+3}}{2k+1} = \sum_{k=0}^{\infty} (-1)^k \frac{x^{6k+4}}{2k+1}$$

Find the Taylor Series of the following functions at given a .

TAYLOR SERIES

Problem 10 $f(x) = x - x^3$ at $a = -2$

Solution

Taylor Series of $f(x) = x - x^3$ at $a = -2$ is of the form

$$\begin{aligned} & f(-2) + f^{(1)}(-2)(x+2) + \frac{f^{(2)}(-2)}{2!}(x+2)^2 \\ & + \frac{f^{(3)}(-2)}{3!}(x+2)^3 + \frac{f^{(4)}(-2)}{4!}(x+2)^4 + \dots \end{aligned}$$

TAYLOR SERIES

Solution(cont'd)

Since f is a polynomial function of degree 3, its derivatives of order higher than 3 is 0. Thus Taylor Series is of the form

$$f(-2) + f^{(1)}(-2)(x+2) + \frac{f^{(2)}(-2)}{2!}(x+2)^2 + \frac{f^{(3)}(-2)}{3!}(x+2)^3$$

TAYLOR SERIES

Solution(cont'd)

By direct computation,

$$f(-2) = 6, \quad f^{(1)}(-2) = -11, \quad f^{(2)}(-2) = 12, \quad f^{(3)}(-2) = -6$$

So the Taylor Series of $x - x^3$ at $a = -2$ is

$$6 - 11(x + 2) + 6(x + 2)^2 - (x + 2)^3$$

TAYLOR SERIES

Problem 11

$$f(x) = \frac{1}{x} \text{ at } a = 2$$

Solution

Taylor Series of $f(x) = 1/x$ at $a = 2$ is of the form

$\sum_{k=0}^{\infty} \frac{f^{(k)}(2)}{k!} (x-2)^k$. We need to find the general expression of the k^{th} derivative of $1/x$.

TAYLOR SERIES

Solution(cont'd)

We derive $1/x$ until a pattern is found.

$$f(x) = 1/x = x^{-1}, \quad f^{(1)}(x) = (-1)x^{-2}$$

$$f^{(2)}(x) = (-1)(-2)x^{-3}, \quad f^{(3)}(x) = (-1)(-2)(-3)x^{-4}$$

In general, $f^{(k)}(x) = (-1)^k k! x^{-(k+1)}$. Therefore

$$f^{(k)}(2) = (-1)^k k! 2^{-(k+1)}.$$

TAYLOR SERIES

Solution(cont'd)

After inserting the general expression of the k^{th} derivative evaluated at 2 we obtain,

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(2)}{k!} (x-2)^k = \sum_{k=0}^{\infty} \frac{1}{k!} (-1)^k k! 2^{-(k+1)} (x-2)^k$$

Hence, the the Taylor Series of $\frac{1}{x}$ is $\sum_{k=0}^{\infty} \frac{(-1)^k}{2^{(k+1)}} (x-2)^k$.

TAYLOR SERIES

Problem 12

$$f(x) = e^{-2x} \text{ at } a = 1/2$$

Solution

Taylor Series of $f(x) = e^{-2x}$ at $a = 1/2$ is of the form

$\sum_{k=0}^{\infty} \frac{f^{(k)}(1/2)}{k!} \left(x - \frac{1}{2}\right)^k$. We need to find the general expression of the k^{th} derivative of e^{-2x} .

TAYLOR SERIES

Solution(cont'd)

We derive e^{-2x} until a pattern is found.

$$f(x) = e^{-2x}, \quad f^{(1)}(x) = -2e^{-2x}, \quad f^{(2)}(x) = -2 - 2e^{-2x}$$

$$\text{In general, } f^{(k)}(x) = (-1)^k 2^k e^{-2x}.$$

$$\text{Therefore } f^{(k)}(1/2) = (-1)^k 2^k e^{-2 \times \frac{1}{2}} = \frac{(-1)^k 2^k}{e}.$$

TAYLOR SERIES

Solution(cont'd)

After inserting the general expression of the k^{th} derivative evaluated at $1/2$ we obtain,

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(1/2)}{k!} \left(x - \frac{1}{2}\right)^k = \sum_{k=0}^{\infty} \frac{1}{k!} \frac{(-1)^k 2^k}{e} \left(x - \frac{1}{2}\right)^k$$

Hence, the the Taylor Series of e^{-2x} is

$$\sum_{k=0}^{\infty} \frac{(-1)^k}{e \times (k!)} (2x - 1)^k.$$

TAYLOR SERIES

Problem 13

$$f(x) = \sin(x) \text{ at } a = \pi/4$$

Solution

Taylor Series of $f(x) = \sin(x)$ at $a = \pi/4$ is of the form $\sum_{k=0}^{\infty} \frac{f^{(k)}(\pi/4)}{k!} \left(x - \frac{\pi}{4}\right)^k$. We need to find the general expression of the k^{th} derivative of $\sin(x)$.

TAYLOR SERIES

Solution(cont'd)

We derive $\sin(x)$ until a pattern is found.

$$f(x) = \sin(x), \quad f^{(1)}(x) = \cos(x), \quad f^{(2)}(x) = -\sin(x)$$

$$\text{In general, } f^{(k)}(x) = \begin{cases} \sin(x) & \text{if } k = 4n \\ \cos(x) & \text{if } k = 4n+1 \\ -\sin(x) & \text{if } k = 4n+2 \\ -\cos(x) & \text{if } k = 4n+3 \end{cases}$$

TAYLOR SERIES

Solution(cont'd)

In other words, even order derivatives are either $\sin(x)$ or $-\sin(x)$ and odd order derivatives are either $\cos(x)$ or $-\cos(x)$. So the Taylor Series at $a = \pi/4$ can be written as

$$\sum_{k=0}^{\infty} (-1)^k \frac{\sin(\pi/4)}{(2k)!} \left(x - \frac{\pi}{4}\right)^{2k} + \sum_{k=0}^{\infty} (-1)^k \frac{\cos(\pi/4)}{(2k+1)!} \left(x - \frac{\pi}{4}\right)^{2k+1}$$

TAYLOR SERIES

Solution(cont'd)

Since, at $a = \pi/4$, $\sin(\pi/4) = \cos(\pi/4) = 1/\sqrt{2}$, the Taylor Series can be simplified to

$$\sum_{k=0}^{\infty} \frac{(-1)^k}{\sqrt{2}(2k)!} \left(x - \frac{\pi}{4}\right)^{2k} + \sum_{k=0}^{\infty} \frac{(-1)^k}{\sqrt{2}(2k+1)!} \left(x - \frac{\pi}{4}\right)^{2k+1}.$$

TAYLOR SERIES

Problem 14

$$f(x) = 10^x \text{ at } a = 1$$

Solution

Taylor Series of $f(x) = 10^x$ at $a = 1$ is of the form

$\sum_{k=0}^{\infty} \frac{f^{(k)}(1)}{k!} (x-1)^k$. We need to find the general expression of the k^{th} derivative of 10^x .

TAYLOR SERIES

Solution(cont'd)

We derive 10^x until a pattern is found.

$$f(x) = 10^x, \quad f^{(1)}(x) = \ln(10) \times 10^x, \quad f^{(2)}(x) = \ln^2(10) 10^x$$

In general, $f^{(k)}(x) = \ln^k(10) 10^x$.

Therefore $f^{(k)}(1) = \ln^k(10) 10$.

TAYLOR SERIES

Solution(cont'd)

After inserting the general expression of the k^{th} derivative evaluated at 1 we obtain,

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(1)}{k!} (x-1)^k = \sum_{k=0}^{\infty} \frac{\ln^k(10)10}{k!} (x-1)^k$$

TAYLOR SERIES

Problem 15

$$f(x) = \ln(1+x) \text{ at } a = -2$$

Solution

Taylor Series of $f(x) = \ln(1+x)$ at $a = -2$ is of the form $\sum_{k=0}^{\infty} \frac{f^{(k)}(-2)}{k!} (x+2)^k$. We need to find the general expression of the k^{th} derivative of $\ln(1+x)$.

TAYLOR SERIES

Solution(cont'd)

We derive $\ln(x+1)$ until a pattern is found.

$$f(x) = \ln(x+1), \quad f^{(1)}(x) = \frac{1}{x+1}, \quad f^{(2)}(x) = -\frac{1}{(x+1)^2}$$

$$\text{In general, } f^{(k)}(x) = \frac{(-1)^k}{(x+1)^k}. \text{ Therefore } f^{(k)}(-2) = 1.$$

TAYLOR SERIES

Solution(cont'd)

After inserting the general expression of the k^{th} derivative evaluated at -2 we obtain

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(-2)}{k!} (x+2)^k = \sum_{k=0}^{\infty} \frac{1}{k!} (x+2)^k.$$